

# A BLUEPRINT FOR THE FORMALIZATION OF SEYMOUR'S MATROID DECOMPOSITION THEOREM

IVAN SERGEEV, MARTIN DVORAK, MARK SANDEY, PIETRO MONTICONE,  
AND CAMERON RAMPELL

ABSTRACT. This document is a blueprint for the formalization in Lean of the structural theory of regular matroids underlying Seymour's decomposition theorem. We present a modular account of regularity via totally unimodular representations, show that regularity is preserved under 1-, 2-, and 3-sums, and establish regularity for several special classes of matroids, including graphic, cographic, and the matroid  $R_{10}$ .

The blueprint records the logical structure of the proof, the precise dependencies between results, and their correspondence with Lean declarations. It is intended both as a guide for the ongoing formalization effort and as a human-readable reference for the organization of the proof.

## CONTENTS

1. Introduction	1
2. Preliminaries	2
2.1. Total Unimodularity	2
2.2. Pivoting	3
2.3. Vector Matroids	4
2.4. Regular Matroids	5
3. Regularity of 1-Sum	6
4. Regularity of 2-Sum	6
5. Regularity of 3-Sum	7
5.1. Definition	7
5.2. Canonical Signing	8
5.3. Properties of Canonical Signing	10
5.4. Proof of Regularity	13
6. Special Matroids	14
6.1. Regularity of Cographic Matroids	15
7. Conclusion	17
References	17

## 1. INTRODUCTION

Seymour's decomposition theorem provides a structural characterization of regular matroids by expressing them as iterated 1-, 2-, and 3-sums of graphic matroids, cographic matroids, and a single exceptional matroid  $R_{10}$ . This result lies at the intersection of matroid theory, linear optimization, and combinatorial geometry, and it plays a central role in the theory of totally unimodular matrices and polynomial-time algorithms.

Our presentation of the structural theory of regular matroids closely follows the exposition and terminology of Truemper's monograph [1], which serves as a primary reference for the decomposition theory and the matrix-based approach adopted throughout this blueprint.

The present document is a *blueprint* for the formalization of this theory in the Lean proof assistant. Rather than presenting a traditional mathematical exposition, the blueprint records

the logical structure of the proof, isolates intermediate results into modular components, and tracks the precise dependencies between statements. Each definition, lemma, and theorem is intended to correspond to a Lean declaration, and many proofs are deferred to Lean and indicated as such.

The blueprint is organized into several thematic parts. We begin by developing the necessary background on totally unimodular matrices, pivoting operations, and vector matroids. We then prove that regularity is preserved under 1-, 2-, and 3-sums of matroids. Finally, we establish regularity for certain special matroids – graphic matroids, cographic matroids, and the matroid  $R_{10}$  – thereby completing the ingredients needed for Seymour’s decomposition.

## 2. PRELIMINARIES

### 2.1. Total Unimodularity.

**Definition 1.** Matrix is a function that takes a row index and returns a vector, which is a function that takes a column index and returns a value. The former aforementioned identity is definitional, the latter is syntactical. By abuse of notation  $(R^Y)^X \equiv R^{X \times Y}$  we do not curry functions in this text. When a matrix happens to be finite (that is, both  $X$  and  $Y$  are finite) and its entries are numeric, we like to represent it by a table of numbers.

**Definition 2.** Let  $A$  be a square matrix over a commutative ring whose rows and columns are indexed by the integers  $\{1, \dots, n\}$ . The determinant of  $A$  is

$$\det A = \sum_{\sigma \in S_n} \left( \operatorname{sgn}(\sigma) \prod_{i=1}^n a_{i, \sigma(i)} \right),$$

where the sum is computed over all permutations  $\sigma \in S_n$ ,  $\operatorname{sgn}(\sigma)$  denotes the sign of permutation  $\sigma$ , and  $a_{i,j} \in R$  is the element element of  $A$  corresponding to the  $i$ -th row and the  $j$ -th column.

**Definition 3.** Let  $R$  be a commutative ring. We say that a matrix  $A \in R^{X \times Y}$  is totally unimodular, or TU for short, if for every  $k \in \mathbb{N}$ , every (not necessarily contiguous)  $k \times k$  submatrix  $T$  of  $A$  has  $\det T \in \{0, \pm 1\}$ .

**Lemma 4.** Let  $A$  be a TU matrix. Suppose rows of  $A$  are multiplied by  $\{0, \pm 1\}$  factors. Then the resulting matrix  $A'$  is also TU.

*Proof.* We prove that  $A'$  is TU by Definition 3. To this end, let  $T'$  be a square submatrix of  $A'$ . Our goal is to show that  $\det T' \in \{0, \pm 1\}$ . Let  $T$  be the submatrix of  $A$  that represents  $T'$  before pivoting. If some of the rows of  $T$  were multiplied by zeros, then  $T'$  contains zero rows, and hence  $\det T' = 0$ . Otherwise,  $T'$  was obtained from  $T$  by multiplying certain rows by  $-1$ . Since  $T'$  has finitely many rows, the number of such multiplications is also finite. Since multiplying a row by  $-1$  results in the determinant getting multiplied by  $-1$ , we get  $\det T' = \pm \det T \in \{0, \pm 1\}$  as desired.  $\square$

**Lemma 5.** Let  $A$  be a TU matrix. Suppose columns of  $A$  are multiplied by  $\{0, \pm 1\}$  factors. Then the resulting matrix  $A'$  is also TU.

*Proof.* Apply Lemma 4 to  $A^\top$ .  $\square$

**Definition 6.** Given  $k \in \mathbb{N}$ , we say that a matrix  $A$  is  $k$ -partially unimodular, or  $k$ -PU for short, if every (not necessarily contiguous, not necessarily injective)  $k \times k$  submatrix  $T$  of  $A$  has  $\det T \in \{0, \pm 1\}$ .

**Lemma 7.** A matrix  $A$  is TU if and only if  $A$  is  $k$ -PU for every  $k \in \mathbb{N}$ .

*Proof.* This follows from Definitions 3 and 6.  $\square$

**Definition 8.** Matrix made of 4 blocks (2x2).

## 2.2. Pivoting.

**Definition 9.** Let  $A \in R^{X \times Y}$  be a matrix and let  $(x, y) \in X \times Y$  be such that  $A(x, y) \neq 0$ . A long tableau pivot in  $A$  on  $(x, y)$  is the operation that maps  $A$  to the matrix  $A'$  where

$$\forall i \in X, \forall j \in Y, A'(i, j) = \begin{cases} \frac{A(i, j)}{A(x, y)}, & \text{if } i = x, \\ A(i, j) - \frac{A(i, y) \cdot A(x, j)}{A(x, y)}, & \text{if } i \neq x. \end{cases}$$

**Lemma 10.** Let  $A \in \mathbb{Q}^{X \times Y}$  be a TU matrix and let  $(x, y) \in X \times Y$  be such that  $A(x, y) \neq 0$ . Then performing the long tableau pivot in  $A$  on  $(x, y)$  yields a TU matrix.

*Proof.* See implementation in Lean.  $\square$

**Definition 11.** Let  $A \in R^{X \times Y}$  be a matrix and let  $(x, y) \in X \times Y$  be such that  $A(x, y) \neq 0$ . Perform the following sequence of operations.

- (1) Adjoin the identity matrix  $1 \in R^{X \times X}$  to  $A$ , resulting in the matrix  $B = \begin{bmatrix} 1 & A \end{bmatrix} \in R^{X \times (X \oplus Y)}$ .
- (2) Perform a long tableau pivot in  $B$  on  $(x, y)$ , and let  $C$  denote the result.
- (3) Swap columns  $x$  and  $y$  in  $C$ , and let  $D$  be the resulting matrix.
- (4) Finally, remove columns indexed by  $X$  from  $D$ , and let  $A'$  be the resulting matrix.

A short tableau pivot in  $A$  on  $(x, y)$  is the operation that maps  $A$  to the matrix  $A'$  defined above.

**Lemma 12.** Let  $A \in R^{X \times Y}$  be a matrix and let  $(x, y) \in X \times Y$  be such that  $A(x, y) \neq 0$ . Then the short tableau pivot in  $A$  on  $(x, y)$  maps  $A$  to  $A'$  with

$$\forall i \in X, \forall j \in Y, A'(i, j) = \begin{cases} \frac{1}{A(x, y)}, & \text{if } i = x \text{ and } j = y, \\ \frac{A(x, j)}{A(x, y)}, & \text{if } i = x \text{ and } j \neq y, \\ -\frac{A(i, j)}{A(x, y)}, & \text{if } i \neq x \text{ and } j = y, \\ A(i, j) - \frac{A(i, y) \cdot A(x, j)}{A(x, y)}, & \text{if } i \neq x \text{ and } j \neq y. \end{cases}$$

*Proof.* Follows by direct calculation.  $\square$

**Lemma 13.** Let  $B = \begin{bmatrix} B_{11} & 0 \\ B_{21} & B_{22} \end{bmatrix} \in \mathbb{Q}^{\{X_1 \cup X_2\} \times \{Y_1 \times Y_2\}}$ . Let  $B' = \begin{bmatrix} B'_{11} & B'_{12} \\ B'_{21} & B'_{22} \end{bmatrix}$  be the result of performing a short tableau pivot on  $(x, y) \in X_1 \times Y_1$  in  $B$ . Then  $B'_{12} = 0$ ,  $B'_{22} = B_{22}$ , and  $\begin{bmatrix} B'_{11} \\ B'_{21} \end{bmatrix}$  is the matrix resulting from performing a short tableau pivot on  $(x, y)$  in  $\begin{bmatrix} B_{11} \\ B_{21} \end{bmatrix}$ .

*Proof.* This follows by a direct calculation. Indeed, because of the 0 block in  $B$ ,  $B_{12}$  and  $B_{22}$  remain unchanged, and since  $\begin{bmatrix} B_{11} \\ B_{21} \end{bmatrix}$  is a submatrix of  $B$  containing the pivot element, performing a short tableau pivot in it is equivalent to performing a short tableau pivot in  $B$  and then taking the corresponding submatrix.  $\square$

**Lemma 14.** Let  $k \in \mathbb{N}$ , let  $A \in \mathbb{Q}^{k \times k}$ , and let  $A'$  be the result of performing a short tableau pivot in  $A$  on  $(x, y)$  with  $x, y \in \{1, \dots, k\}$  such that  $A(x, y) \neq 0$ . Then  $A'$  contains a submatrix  $A''$  of size  $(k-1) \times (k-1)$  with  $|\det A''| = |\det A|/|A(x, y)|$ .

*Proof.* Let  $X = \{1, \dots, k\} \setminus \{x\}$  and  $Y = \{1, \dots, k\} \setminus \{y\}$ , and let  $A'' = A'(X, Y)$ . Since  $A''$  does not contain the pivot row or the pivot column,  $\forall (i, j) \in X \times Y$  we have  $A''(i, j) = A(i, j) - \frac{A(i, y) \cdot A(x, j)}{A(x, y)}$ . For  $\forall j \in Y$ , let  $B_j$  be the matrix obtained from  $A$  by removing row  $x$  and column  $j$ , and let  $B'_j$  be the matrix obtained from  $A''$  by replacing column  $j$  with  $A(X, y)$  (i.e., the pivot column without the pivot element). The cofactor expansion along row  $x$  in  $A$  yields

$$\det A = \sum_{j=1}^k (-1)^{y+j} \cdot A(x, j) \cdot \det B_j.$$

By reordering columns of every  $B_j$  to match their order in  $B_j''$ , we get

$$\det A = (-1)^{x+y} \cdot \left( A(x, y) \cdot \det A' - \sum_{j \in Y} A(x, j) \cdot \det B_j'' \right).$$

By linearity of the determinant applied to  $\det A''$ , we have

$$\det A'' = \det A' - \sum_{j \in Y} \frac{A(x, j)}{A(x, y)} \cdot \det B_j''$$

Therefore,  $|\det A''| = |\det A|/|A(x, y)|$ . □

**Lemma 15.** Let  $A \in \mathbb{Q}^{X \times Y}$  be a TU matrix and let  $(x, y) \in X \times Y$  be such that  $A(x, y) \neq 0$ . Then performing the short tableau pivot in  $A$  on  $(x, y)$  yields a TU matrix.

*Proof.* See implementation in Lean. □

### 2.3. Vector Matroids.

**Definition 16.** A matroid  $M$  is a pair  $(E, \mathcal{I})$  where  $E$  is a (possibly infinite) set and  $\mathcal{I} \in 2^E$  is such that:

- (1)  $\emptyset \in \mathcal{I}$
- (2) If  $I \in \mathcal{I}$  and  $J \subseteq I$ , then  $J \in \mathcal{I}$ .
- (3) If  $I \in \mathcal{I}$  is not maximal (with respect to set inclusion) and  $B \in \mathcal{I}$  is maximal, then there exists an  $x \in B \setminus I$  such that  $I \cup \{x\} \in \mathcal{I}$ .
- (4) If  $X \subseteq E$  and  $I \subseteq X$  is such that  $I \in \mathcal{I}$ , then there exists an  $J \in \mathcal{I}$  with  $I \subseteq J \subseteq X$  that is maximal with respect to set inclusion.

We call  $E$  the ground set of  $M$  and  $\mathcal{I}$  the collection of independent sets in  $M$ . We say that  $B \in \mathcal{I}$  is a base of  $M$  if  $B$  is maximal in  $\mathcal{I}$ .

**Definition 17.** Let  $R$  be a division ring, let  $X$  and  $Y$  be sets, and let  $A \in R^{X \times Y}$  be a matrix. The vector matroid of  $A$  is the matroid  $M = (Y, \mathcal{I})$  where a set  $I \subset Y$  is independent in  $M$  if and only if the columns of  $A$  indexed by  $I$  are linearly independent.

**Definition 18.** Let  $R$  be a division ring, let  $X$  and  $Y$  be disjoint sets, and let  $S \in R^{X \times Y}$  be a matrix. Let  $A = [1 \ S] \in R^{X \times (X \cup Y)}$  be the matrix obtained from  $S$  by adjoining the identity matrix as columns, and let  $M$  be the vector matroid of  $A$ . Then  $S$  is called the standard representation of  $M$ .

**Lemma 19.** Let  $S \in R^{X \times Y}$  be a standard representation of a vector matroid  $M$ . Then  $X$  is a base in  $M$ .

*Proof.* See implementation in Lean. □

**Lemma 20.** Adding extra zero rows to a full representation matrix of a vector matroid does not change the matroid.

*Proof.* See implementation in Lean. □

**Lemma 21.** Let  $A \in \mathbb{Q}^{X \times Y}$  be a TU matrix, let  $M$  be the vector matroid of  $A$ , and let  $B$  be a base of  $M$ . Then there exists a matrix  $S \in \mathbb{Q}^{B \times (Y \setminus B)}$  such that  $S$  is TU and  $S$  is a standard representation of  $M$ .

*Proof.* See implementation in Lean. □

**Definition 22.** Let  $R$  be a magma containing zero. The support of matrix  $A \in R^{X \times Y}$  is  $A^\# \in \{0, 1\}^{X \times Y}$  given by

$$\forall i \in X, \forall j \in Y, A^\#(i, j) = \begin{cases} 0, & \text{if } A(i, j) = 0, \\ 1, & \text{if } A(i, j) \neq 0. \end{cases}$$

**Lemma 23.** Transpose of a support matrix is equal to a support of the transposed matrix.

*Proof.* Definitional equality. □

**Lemma 24.** Submatrix of a support matrix is equal to a support matrix of the submatrix.

*Proof.* Definitional equality. □

**Lemma 25.** If  $A$  is a matrix over  $\mathbb{Z}_2$ , then  $A^\# = A$ .

*Proof.* Check elementwise equality. □

**Lemma 26.** If two standard representation matrices of the same matroid have the same base, then they have the same support.

*Proof.* See implementation in Lean. □

**Lemma 27.** A square matrix is invertible iff its determinant is invertible.

*Proof.* This result is proved in Mathlib. □

**Lemma 28.** Let  $A$  be a rational TU matrix with finite number of rows and finite number of columns. Its rows are linearly independent iff the rows of its support matrix are linearly independent.

*Proof.* See implementation in Lean. □

**Lemma 29.** Let  $A$  be a rational TU matrix with finite number of rows. Its rows are linearly independent iff the rows of its support matrix are linearly independent.

*Proof.* See implementation in Lean. □

**Lemma 30.** Let  $A$  be a rational TU matrix. Its rows are linearly independent iff the rows of its support matrix are linearly independent.

*Proof.* See implementation in Lean. □

**Lemma 31.** Let  $A$  be a TU matrix.

- (1) If a matroid is represented by  $A$ , then it is also represented by  $A^\#$ .
- (2) If a matroid is represented by  $A^\#$ , then it is also represented by  $A$ .

*Proof.* See implementation in Lean. □

## 2.4. Regular Matroids.

**Definition 32.** A matroid  $M$  is regular if there exists a TU matrix  $A \in \mathbb{Q}^{X \times Y}$  such that  $M$  is a vector matroid of  $A$ .

**Definition 33.** We say that  $A' \in \mathbb{Q}^{X \times Y}$  is a TU signing of  $A \in \mathbb{Z}_2^{X \times Y}$  if  $A'$  is TU and

$$\forall i \in X, \forall j \in Y, |A'(i, j)| = A(i, j).$$

**Lemma 34.** Let  $B \in \mathbb{Z}_2^{X \times Y}$  be a standard representation matrix of a matroid  $M$ . Then  $M$  is regular if and only if  $B$  has a TU signing.

*Proof.* Suppose that  $M$  is regular. By Definition 32, there exists a TU matrix  $A \in \mathbb{Q}^{X \times Y}$  such that  $M$  is a vector matroid of  $A$ . By Lemma 19,  $X$  (the row set of  $B$ ) is a base of  $M$ . By Lemma 21,  $A$  can be converted into a standard representation matrix  $B' \in \mathbb{Q}^{X \times Y}$  of  $M$  such that  $B'$  is also TU. Since  $B'$  and  $B$  are both standard representations of  $M$ , by Lemma 26 the support matrices  $(B')^\#$  and  $B^\#$  are the same. Lemma 25 gives  $B^\# = B$ . Thus,  $B'$  is TU and  $(B')^\# = B$ , so  $B'$  is a TU signing of  $B$ .

Suppose that  $B$  has a TU signing  $B' \in \mathbb{Q}^{X \times Y}$ . Then  $A = [1 \mid B']$  is TU, as it is obtained from  $B'$  by adjoining the identity matrix. Moreover, by Lemma 31,  $A$  represents the same matroid as  $A^\# = [1 \mid B]$ , which is  $M$ . Thus,  $A$  is a TU matrix representing  $M$ , so  $M$  is regular. □

## 3. REGULARITY OF 1-SUM

**Definition 35.** Let  $R$  be a magma containing zero (we will use  $R = \mathbb{Z}_2$  and  $R = \mathbb{Q}$ ). Let  $B_\ell \in R^{X_\ell \times Y_\ell}$  and  $B_r \in R^{X_r \times Y_r}$  be matrices where  $X_\ell, Y_\ell, X_r, Y_r$  are pairwise disjoint sets. The 1-sum  $B = B_\ell \oplus_1 B_r$  of  $B_\ell$  and  $B_r$  is

$$B = \begin{bmatrix} B_\ell & 0 \\ 0 & B_r \end{bmatrix} \in R^{(X_\ell \cup X_r) \times (Y_\ell \cup Y_r)}.$$

**Definition 36.** A matroid  $M$  is a 1-sum of matroids  $M_\ell$  and  $M_r$  if there exist standard  $\mathbb{Z}_2$  representation matrices  $B_\ell, B_r$ , and  $B$  (for  $M_\ell, M_r$ , and  $M$ , respectively) of the form given in Definition 35.

**Lemma 37.** Let  $A$  be a square matrix of the form  $A = \begin{bmatrix} A_{11} & A_{12} \\ 0 & A_{22} \end{bmatrix}$ . Then  $\det A = \det A_{11} \cdot \det A_{22}$ .

*Proof.* This result is proved in Mathlib.  $\square$

**Lemma 38.** Let  $B_\ell$  and  $B_r$  from Definition 35 be TU matrices (over  $\mathbb{Q}$ ). Then  $B = B_\ell \oplus_1 B_r$  is TU.

*Proof.* We prove that  $B$  is TU by Definition 3. To this end, let  $T$  be a square submatrix of  $B$ . Our goal is to show that  $\det T \in \{0, \pm 1\}$ .

Let  $T_\ell$  and  $T_r$  denote the submatrices in the intersection of  $T$  with  $B_\ell$  and  $B_r$ , respectively. Then  $T$  has the form

$$T = \begin{bmatrix} T_\ell & 0 \\ 0 & T_r \end{bmatrix}.$$

First, suppose that  $T_\ell$  and  $T_r$  are square. Then  $\det T = \det T_\ell \cdot \det T_r$  by Lemma 37. Moreover,  $\det T_\ell, \det T_r \in \{0, \pm 1\}$ , since  $T_\ell$  and  $T_r$  are square submatrices of TU matrices  $B_\ell$  and  $B_r$ , respectively. Thus,  $\det T \in \{0, \pm 1\}$ , as desired.

Without loss of generality we may assume that  $T_\ell$  has fewer rows than columns. Otherwise we can transpose all matrices and use the same proof, since TUness and determinants are preserved under transposition. Thus,  $T$  can be represented in the form

$$T = \begin{bmatrix} T_{11} & T_{12} \\ 0 & T_{22} \end{bmatrix},$$

where  $T_{11}$  contains  $T_\ell$  and some zero rows,  $T_{22}$  is a submatrix of  $T_r$ , and  $T_{12}$  contains the rest of the rows of  $T_r$  (not contained in  $T_{22}$ ) and some zero rows. By Lemma 37, we have  $\det T = \det T_{11} \cdot \det T_{22}$ . Since  $T_{11}$  contains at least one zero row,  $\det T_{11} = 0$ . Thus,  $\det T = 0 \in \{0, \pm 1\}$ , as desired.  $\square$

**Theorem 39.** Let  $M$  be a 1-sum of regular matroids  $M_\ell$  and  $M_r$ . Then  $M$  is also regular.

*Proof.* Let  $B_\ell, B_r$ , and  $B$  be standard  $\mathbb{Z}_2$  representation matrices from Definition 36. Since  $M_\ell$  and  $M_r$  are regular, by Lemma 34,  $B_\ell$  and  $B_r$  have TU signings  $B'_\ell$  and  $B'_r$ , respectively. Then  $B' = B'_\ell \oplus_1 B'_r$  is a TU signing of  $B$ . Indeed,  $B'$  is TU by Lemma 38, and a direct calculation shows that  $B'$  is a signing of  $B$ . Thus,  $M$  is regular by Lemma 34.  $\square$

## 4. REGULARITY OF 2-SUM

**Definition 40.** Let  $R$  be a semiring (we will use  $R = \mathbb{Z}_2$  and  $R = \mathbb{Q}$ ). Let  $B_\ell \in R^{X_\ell \times Y_\ell}$  and  $B_r \in R^{X_r \times Y_r}$  where  $X_\ell \cap X_r = \{x\}$ ,  $Y_\ell \cap Y_r = \{y\}$ ,  $X_\ell$  is disjoint with  $Y_\ell$  and  $Y_r$ , and  $X_r$  is disjoint with  $Y_\ell$  and  $Y_r$ . Additionally, let  $A_\ell = B_\ell(X_\ell \setminus \{x\}, Y_\ell)$  and  $A_r = B_r(X_r, Y_r \setminus \{y\})$ , and suppose  $r = B_\ell(x, Y_\ell) \neq 0$  and  $c = B_r(X_r, y) \neq 0$ . Then the 2-sum  $B = B_\ell \oplus_{2,x,y} B_r$  of  $B_\ell$  and  $B_r$  is defined as

$$B = \begin{bmatrix} A_\ell & 0 \\ D & A_r \end{bmatrix} \quad \text{where } D = c \otimes r.$$

Here  $D \in R^{X_r \times Y_\ell}$ , and the indexing is consistent everywhere.

**Definition 41.** A matroid  $M$  is a 2-sum of matroids  $M_\ell$  and  $M_r$  if there exist standard  $\mathbb{Z}_2$  representation matrices  $B_\ell$ ,  $B_r$ , and  $B$  (for  $M_\ell$ ,  $M_r$ , and  $M$ , respectively) of the form given in Definition 40.

**Lemma 42.** Let  $B_\ell$  and  $B_r$  from Definition 40 be TU matrices (over  $\mathbb{Q}$ ). Then  $C = \begin{bmatrix} D & A_r \end{bmatrix}$  is TU.

*Proof.* Since  $B_\ell$  is TU, all its entries are in  $\{0, \pm 1\}$ . In particular,  $r$  is a  $\{0, \pm 1\}$  vector. Therefore, every column of  $D$  is a copy of  $y$ ,  $-y$ , or the zero column. Thus,  $C$  can be obtained from  $B_r$  by adjoining zero columns, duplicating the  $y$  column, and multiplying some columns by  $-1$ . Since all these operations preserve TUness and since  $B_r$  is TU,  $C$  is also TU.  $\square$

**Lemma 43.** Let  $B_\ell$  and  $B_r$  be matrices from Definition 40. Let  $B'_\ell$  and  $B'$  be the matrices obtained by performing a short tableau pivot on  $(x_\ell, y_\ell) \in X_\ell \times Y_\ell$  in  $B_\ell$  and  $B$ , respectively. Then  $B' = B'_\ell \oplus_{2,x,y} B_r$ .

*Proof.* Let

$$B'_\ell = \begin{bmatrix} A'_\ell \\ r' \end{bmatrix}, \quad B' = \begin{bmatrix} B'_{11} & B'_{12} \\ B'_{21} & B'_{22} \end{bmatrix}$$

where the blocks have the same dimensions as in  $B_\ell$  and  $B$ , respectively. By Lemma 13,  $B'_{11} = A'_\ell$ ,  $B'_{12} = 0$ , and  $B'_{22} = A_r$ . Equality  $B'_{21} = c \otimes r'$  can be verified via a direct calculation. Thus,  $B' = B'_\ell \oplus_{2,x,y} B_r$ .  $\square$

**Lemma 44.** Let  $B_\ell$  and  $B_r$  from Definition 40 be TU matrices (over  $\mathbb{Q}$ ). Then  $B_\ell \oplus_{2,x,y} B_r$  is TU.

*Proof.* By Lemma 7, it suffices to show that  $B_\ell \oplus_{2,x,y} B_r$  is  $k$ -PU for every  $k \in \mathbb{N}$ . We prove this claim by induction on  $k$ . The base case with  $k = 1$  holds, since all entries of  $B_\ell \oplus_{2,x,y} B_r$  are in  $\{0, \pm 1\}$  by construction.

Suppose that for some  $k \in \mathbb{N}$  we know that for any TU matrices  $B'_\ell$  and  $B'_r$  (from Definition 40) their 2-sum  $B'_\ell \oplus_{2,x,y} B'_r$  is  $k$ -PU. Now, given TU matrices  $B_\ell$  and  $B_r$  (from Definition 40), our goal is to show that  $B = B_\ell \oplus_{2,x,y} B_r$  is  $(k+1)$ -PU, i.e., that every  $(k+1) \times (k+1)$  submatrix  $T$  of  $B$  has  $\det T \in \{0, \pm 1\}$ .

First, suppose that  $T$  has no rows in  $X_\ell$ . Then  $T$  is a submatrix of  $\begin{bmatrix} D & A_r \end{bmatrix}$ , which is TU by Lemma 42, so  $\det T \in \{0, \pm 1\}$ . Thus, we may assume that  $T$  contains a row  $x_\ell \in X_\ell$ .

Next, note that without loss of generality we may assume that there exists  $y_\ell \in Y_\ell$  such that  $T(x_\ell, y_\ell) \neq 0$ . Indeed, if  $T(x_\ell, y) = 0$  for all  $y$ , then  $\det T = 0$  and we are done, and  $T(x_\ell, y) = 0$  holds whenever  $y \in Y_r$ .

Since  $B$  is 1-PU, all entries of  $T$  are in  $\{0, \pm 1\}$ , and hence  $T(x_\ell, y_\ell) \in \{\pm 1\}$ . Thus, by Lemma 14, performing a short tableau pivot in  $T$  on  $(x_\ell, y_\ell)$  yields a matrix that contains a  $k \times k$  submatrix  $T''$  such that  $|\det T| = |\det T''|$ . Since  $T$  is a submatrix of  $B$ , matrix  $T''$  is a submatrix of the matrix  $B'$  resulting from performing a short tableau pivot in  $B$  on the same entry  $(x_\ell, y_\ell)$ . By Lemma 43, we have  $B' = B'_\ell \oplus_{2,x,y} B_r$  where  $B'_\ell$  is the result of performing a short tableau pivot in  $B_\ell$  on  $(x_\ell, y_\ell)$ . Since  $B_\ell$  is TU, by Lemma 15,  $B'_\ell$  is also TU. Thus, by the inductive hypothesis applied to  $T''$  and  $B'_\ell \oplus_{2,x,y} B_r$ , we have  $\det T'' \in \{0, \pm 1\}$ . Since  $|\det T| = |\det T''|$ , we conclude that  $\det T \in \{0, \pm 1\}$ .  $\square$

**Theorem 45.** Let  $M$  be a 2-sum of regular matroids  $M_\ell$  and  $M_r$ . Then  $M$  is also regular.

*Proof.* Let  $B_\ell$ ,  $B_r$ , and  $B$  be standard  $\mathbb{Z}_2$  representation matrices from Definition 41. Since  $M_\ell$  and  $M_r$  are regular, by Lemma 34,  $B_\ell$  and  $B_r$  have TU signings  $B'_\ell$  and  $B'_r$ , respectively. Then  $B' = B'_\ell \oplus_{2,x,y} B'_r$  is a TU signing of  $B$ . Indeed,  $B'$  is TU by Lemma 44, and a direct calculation verifies that  $B'$  is a signing of  $B$ . Thus,  $M$  is regular by Lemma 34.  $\square$

## 5. REGULARITY OF 3-SUM

### 5.1. Definition.

**Definition 46.** Let  $X_\ell$ ,  $Y_\ell$ ,  $X_r$ , and  $Y_r$  be sets satisfying the following properties:

- $X_\ell \cap X_r = \{x_2, x_1, x_0\}$  for some distinct  $x_0, x_1$ , and  $x_2$ ;
- $Y_\ell \cap Y_r = \{y_0, y_1, y_2\}$  for some distinct  $y_0, y_1$ , and  $y_2$ ;
- $X_\ell$  is disjoint with  $Y_r$ ; and
- $Y_\ell$  is disjoint with  $X_r$ .

Let  $B_\ell \in \mathbb{Z}_2^{X_\ell \times Y_\ell}$  and  $B_r \in \mathbb{Z}_2^{X_r \times Y_r}$  be matrices of the form

$$B_\ell = \begin{array}{|c|c|c|} \hline & & \\ \hline & A_\ell & 0 \\ \hline & 1 & 1 & 0 \\ \hline D_\ell & D_0 & 1 \\ \hline & & 1 \\ \hline \end{array} \quad \text{and} \quad B_r = \begin{array}{|c|c|c|c|} \hline 1 & 1 & 0 & 0 \\ \hline D_0 & 1 & & \\ \hline & 1 & & A_r \\ \hline D_r & & & \\ \hline \end{array}$$

where  $D_0$  is invertible. Then the 3-sum  $B = B_\ell \oplus_3 B_r$  of  $B_\ell$  and  $B_r$  is defined as

$$B = \begin{array}{|c|c|c|} \hline & & \\ \hline & A_\ell & 0 \\ \hline & 1 & 1 & 0 \\ \hline D_\ell & D_0 & 1 \\ \hline & & 1 & A_r \\ \hline D_{\ell r} & D_r & & \\ \hline \end{array} \quad \text{where } D_{\ell r} = D_r \cdot (D_0)^{-1} \cdot D_\ell.$$

Here the indexing is consistent between all the matrices,  $D_0 \in \mathbb{Z}_2^{\{x_1, x_0\} \times \{y_0, y_1\}}$ , and the submatrix

$$\begin{array}{|c|c|c|} \hline 1 & 1 & 0 \\ \hline D_0 & 1 & \\ \hline & 1 & \\ \hline \end{array}$$

is indexed by  $\{x_2, x_1, x_0\} \times \{y_0, y_1, y_2\}$  in  $B_\ell$ ,  $B_r$ , and  $B$ .

**Definition 47.** A matroid  $M$  is a 3-sum of matroids  $M_\ell$  and  $M_r$  if there exist standard  $\mathbb{Z}_2$  representation matrices  $B_\ell$ ,  $B_r$ , and  $B$  (for  $M_\ell$ ,  $M_r$ , and  $M$ , respectively) of the form given in Definition 46.

## 5.2. Canonical Signing.

**Lemma 48.** Let  $D_0 \in \mathbb{Z}_2^{\{x_1, x_0\} \times \{y_0, y_1\}}$  be an invertible matrix. Then, up to reindexing of rows and columns, either  $D_0 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  or  $D_0 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ .

*Proof.* Brute force. □

For the sake of simplicity of notation, going forward we assume that the submatrix  $D_0$  in Definition 46 falls into one of the two special cases presented in Lemma 48.

**Definition 49.** We call  $D'_0 \in \mathbb{Q}^{\{x_1, x_0\} \times \{y_0, y_1\}}$  the canonical signing of  $D_0 \in \mathbb{Z}_2^{\{x_1, x_0\} \times \{y_0, y_1\}}$  if

$$D_0 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad D'_0 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad \text{or} \quad D_0 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad D'_0 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}.$$

Similarly, we call  $S' \in \mathbb{Q}^{\{x_2, x_1, x_0\} \times \{y_0, y_1, y_2\}}$  the canonical signing of  $S \in \mathbb{Z}_2^{\{x_2, x_1, x_0\} \times \{y_0, y_1, y_2\}}$  if

$$S = \begin{array}{|c|c|c|} \hline 1 & 1 & 0 \\ \hline D_0 & 1 & \\ \hline & 1 & \\ \hline \end{array} \quad \text{and} \quad S' = \begin{array}{|c|c|c|} \hline 1 & 1 & 0 \\ \hline D'_0 & 1 & \\ \hline & 1 & \\ \hline \end{array}$$

To simplify notation, going forward we use  $D_0$ ,  $D'_0$ ,  $S$ , and  $S'$  to refer to the matrices of the form above. Observe that the canonical signing  $S'$  of  $S$  (from Definition 49) is TU.

**Lemma 50.** Let  $Q$  be a TU signing of  $S$  (from Definition 49). Let  $u \in \{0, \pm 1\}^{\{x_2, x_1, x_0\}}$ ,  $v \in \{0, \pm 1\}^{\{y_0, y_1, y_2\}}$ , and  $Q'$  be defined as follows:

$$u(i) = \begin{cases} Q(x_2, y_0) \cdot Q(x_0, y_0), & i = x_0, \\ Q(x_2, y_0) \cdot Q(x_0, y_0) \cdot Q(x_0, y_2) \cdot Q(x_1, y_2), & i = x_1, \\ 1, & i = x_2, \end{cases}$$

$$v(j) = \begin{cases} Q(x_2, y_0), & j = y_0, \\ Q(x_2, y_1), & j = y_1, \\ Q(x_2, y_0) \cdot Q(x_0, y_0) \cdot Q(x_0, y_2), & j = y_2, \end{cases}$$

$$Q'(i, j) = Q(i, j) \cdot u(i) \cdot v(j) \quad \forall i \in \{x_2, x_1, x_0\}, \forall j \in \{y_0, y_1, y_2\}.$$

Then  $Q' = S'$  (from Definition 49).

*Proof.* Since  $Q$  is a TU signing of  $S$  and  $Q'$  is obtained from  $Q$  by multiplying rows and columns by  $\pm 1$  factors,  $Q'$  is also a TU signing of  $S$ . By construction, we have

$$\begin{aligned} Q'(x_2, y_0) &= Q(x_2, y_0) \cdot 1 \cdot Q(x_2, y_0) = 1, \\ Q'(x_2, y_1) &= Q(x_2, y_1) \cdot 1 \cdot Q(x_2, y_1) = 1, \\ Q'(x_2, y_2) &= 0, \\ Q'(x_0, y_0) &= Q(x_0, y_0) \cdot (Q(x_2, y_0) \cdot Q(x_0, y_0)) \cdot Q(x_2, y_0) = 1, \\ Q'(x_0, y_1) &= Q(x_0, y_1) \cdot (Q(x_2, y_0) \cdot Q(x_0, y_0)) \cdot Q(x_2, y_1), \\ Q'(x_0, y_2) &= Q(x_0, y_2) \cdot (Q(x_2, y_0) \cdot Q(x_0, y_0)) \cdot (Q(x_2, y_0) \cdot Q(x_0, y_0) \cdot Q(x_0, y_2)) = 1, \\ Q'(x_1, y_0) &= 0, \\ Q'(x_1, y_1) &= Q(x_1, y_1) \cdot (Q(x_2, y_0) \cdot Q(x_0, y_0) \cdot Q(x_0, y_2) \cdot Q(x_1, y_2)) \cdot (Q(x_2, y_1)), \\ Q'(x_1, y_2) &= Q(x_1, y_2) \cdot (Q(x_2, y_0) \cdot Q(x_0, y_0) \cdot Q(x_0, y_2) \cdot Q(x_1, y_2)) \cdot (Q(x_2, y_0) \cdot Q(x_0, y_0) \cdot Q(x_0, y_2)) = 1. \end{aligned}$$

Thus, it remains to show that  $Q'(x_0, y_1) = S'(x_0, y_1)$  and  $Q'(x_1, y_1) = S'(x_1, y_1)$ .

Consider the entry  $Q'(x_0, y_1)$ . If  $D_0(x_0, y_1) = 0$ , then  $Q'(x_0, y_1) = 0 = S'(x_0, y_1)$ . Otherwise, we have  $D_0(x_0, y_1) = 1$ , and so  $Q'(x_0, y_1) \in \{\pm 1\}$ , as  $Q'$  is a signing of  $S$ . If  $Q'(x_0, y_1) = -1$ , then

$$\det Q'(\{x_0, x_2\}, \{y_0, y_1\}) = \det \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} = 2 \notin \{0, \pm 1\},$$

which contradicts TUness of  $Q'$ . Thus,  $Q'(x_0, y_1) = 1 = S'(x_0, y_1)$ .

Consider the entry  $Q'(x_1, y_1)$ . Since  $Q'$  is a signing of  $S$ , we have  $Q'(x_1, y_1) \in \{\pm 1\}$ . Consider two cases.

- (1) Suppose that  $D_0 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ . If  $Q'(x_1, y_1) = 1$ , then  $\det Q = \det \begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix} = -2 \notin \{0, \pm 1\}$ , which contradicts TUness of  $Q'$ . Thus,  $Q'(x_1, y_1) = -1 = S'(x_1, y_1)$ .
- (2) Suppose that  $D_0 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ . If  $Q'(x_1, y_1) = -1$ , then  $\det Q(\{x_1, x_0\}, \{y_1, y_2\}) = \det \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} = 2 \notin \{0, \pm 1\}$ , which contradicts TUness of  $Q'$ . Thus,  $Q'(x_1, y_1) = 1 = S'(x_1, y_1)$ .

□

**Definition 51.** Let  $X$  and  $Y$  be sets with  $\{x_2, x_1, x_0\} \subseteq X$  and  $\{y_0, y_1, y_2\} \subseteq Y$ . Let  $Q \in \mathbb{Q}^{X \times Y}$  be a TU matrix. Define  $u \in \{0, \pm 1\}^X$ ,  $v \in \{0, \pm 1\}^Y$ , and  $Q'$  as follows:

$$u(i) = \begin{cases} Q(x_2, y_0) \cdot Q(x_0, y_0), & i = x_0, \\ Q(x_2, y_0) \cdot Q(x_0, y_0) \cdot Q(x_0, y_2) \cdot Q(x_1, y_2), & i = x_1, \\ 1, & i = x_2, \\ 1, & i \in X \setminus \{x_2, x_1, x_0\}, \end{cases}$$

$$v(j) = \begin{cases} Q(x_2, y_0), & j = y_0, \\ Q(x_2, y_1), & j = y_1, \\ Q(x_2, y_0) \cdot Q(x_0, y_0) \cdot Q(x_0, y_2), & j = y_2, \\ 1, & j \in Y \setminus \{y_0, y_1, y_2\}, \end{cases}$$

$$Q'(i, j) = Q(i, j) \cdot u(i) \cdot v(j) \quad \forall i \in X, \forall j \in Y.$$

We call  $Q'$  the canonical re-signing of  $Q$ .

**Lemma 52.** Let  $X$  and  $Y$  be sets with  $\{x_2, x_1, x_0\} \subseteq X$  and  $\{y_0, y_1, y_2\} \subseteq Y$ . Let  $Q \in \mathbb{Q}^{X \times Y}$  be a TU signing of  $Q_0 \in \mathbb{Z}_2^{X \times Y}$  such that  $Q_0(\{x_2, x_1, x_0\}, \{y_0, y_1, y_2\}) = S$  (from Definition 49). Then the canonical re-signing  $Q'$  of  $Q$  (from Definition 51) is a TU signing of  $Q_0$  and  $Q'(\{x_2, x_1, x_0\}, \{y_0, y_1, y_2\}) = S'$  (from Definition 49).

*Proof.* Since  $Q$  is a TU signing of  $Q_0$  and  $Q'$  is obtained from  $Q$  by multiplying some rows and columns by  $\pm 1$  factors,  $Q'$  is also a TU signing of  $Q_0$ . Equality  $Q'(\{x_2, x_1, x_0\}, \{y_0, y_1, y_2\}) = S'$  follows from Lemma 50.  $\square$

**Definition 53.** Suppose that  $B_\ell$  and  $B_r$  from Definition 46 have TU signings  $B'_\ell$  and  $B'_r$ , respectively. Let  $B''_\ell$  and  $B''_r$  be the canonical re-signings (from Definition 51) of  $B'_\ell$  and  $B'_r$ , respectively. Let  $A''_\ell, A''_r, D''_\ell, D''_r$ , and  $D''_0$  be blocks of  $B''_\ell$  and  $B''_r$  analogous to blocks  $A_\ell, A_r, D_\ell, D_r$ , and  $D_0$  of  $B_\ell$  and  $B_r$ . The canonical signing  $B''$  of  $B$  is defined as

$$B'' = \begin{array}{|c|c|c|c|} \hline & A''_\ell & & 0 \\ \hline & \begin{array}{|c|c|c|} \hline 1 & 1 & 0 \\ \hline \end{array} & & \\ \hline D''_\ell & D''_0 & \begin{array}{|c|} \hline 1 \\ \hline 1 \\ \hline \end{array} & A''_r \\ \hline D''_{\ell r} & D''_r & & \\ \hline \end{array} \quad \text{where } D''_{\ell r} = D''_r \cdot (D''_0)^{-1} \cdot D''_\ell.$$

Note that  $D''_0$  is non-singular by construction, so  $D''_{\ell r}$  and hence  $B''$  are well-defined.

### 5.3. Properties of Canonical Signing.

**Lemma 54.**  $B''$  from Definition 53 is a signing of  $B$ .

*Proof.* By Lemma 52,  $B''_\ell$  and  $B''_r$  are TU signings of  $B_\ell$  and  $B_r$ , respectively. As a result, blocks  $A''_\ell, A''_r, D''_\ell, D''_r$ , and  $D''_0$  in  $B''$  are signings of the corresponding blocks in  $B$ . Thus, it remains to show that  $D''_{\ell r}$  is a signing of  $D_{\ell r}$ . This can be verified via a direct calculation.  $\square$

**Lemma 55.** Suppose that  $B_r$  from Definition 46 has a TU signing  $B'_r$ . Let  $B''_r$  be the canonical re-signing (from Definition 51) of  $B'_r$ . Let  $c''_0 = B''_r(X_r, y_0)$ ,  $c''_1 = B''_r(X_r, y_1)$ , and  $c''_2 = c''_0 - c''_1$ . Then the following statements hold.

- (1) For every  $i \in X_r$ ,  $[c''_0(i) \ c''_1(i)] \in \{0, \pm 1\}^{\{y_0, y_1\}} \setminus \{[1 \ -1], [-1 \ 1]\}$ .
- (2) For every  $i \in X_r$ ,  $c''_2(i) \in \{0, \pm 1\}$ .
- (3)  $[c''_0 \ c''_2 \ A''_r]$  is TU.
- (4)  $[c''_1 \ c''_2 \ A''_r]$  is TU.
- (5)  $[c''_0 \ c''_1 \ c''_2 \ A''_r]$  is TU.

*Proof.* Throughout the proof we use that  $B_r''$  is TU, which holds by Lemma 52.

- (1) Since  $B_r''$  is TU, all its entries are in  $\{0, \pm 1\}$ , and in particular  $[c_0''(i) \ c_1''(i)] \in \{0, \pm 1\}^{\{y_0, y_1\}}$ . If  $[c_0''(i) \ c_1''(i)] = [1 \ -1]$ , then

$$\det B_r''(\{x_2, i\}, \{y_0, y_1\}) = \det \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = -2 \notin \{0, \pm 1\},$$

which contradicts TUness of  $B_r''$ . Similarly, if  $[c_0''(i) \ c_1''(i)] = [-1 \ 1]$ , then

$$\det B_r''(\{x_2, i\}, \{y_0, y_1\}) = \det \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} = 2 \notin \{0, \pm 1\},$$

which contradicts TUness of  $B_r''$ . Thus, the desired statement holds.

- (2) Follows from item 1 and a direct calculation.  
(3) Performing a short tableau pivot in  $B_r''$  on  $(x_2, y_0)$  yields:

$$B_r'' = \begin{bmatrix} \boxed{1} & 1 & 0 \\ c_0 & c_1 & A_r \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 0 \\ -c_0 & c_1 - c_0 & A_r \end{bmatrix}$$

The resulting matrix can be transformed into  $[c_0'' \ c_2'' \ A_r'']$  by removing row  $x_2$  and multiplying columns  $y_0$  and  $y_1$  by  $-1$ . Since  $B_r''$  is TU and since TUness is preserved under pivoting, taking submatrices, multiplying columns by  $\pm 1$  factors, we conclude that  $[c_0'' \ c_2'' \ A_r'']$  is TU.

- (4) Similar to item 4, performing a short tableau pivot in  $B_r''$  on  $(x_2, y_1)$  yields:

$$B_r'' = \begin{bmatrix} 1 & \boxed{1} & 0 \\ c_0 & c_1 & A_r \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 0 \\ c_0'' - c_1 & -c_1 & A_r \end{bmatrix}$$

The resulting matrix can be transformed into  $[c_1'' \ c_2'' \ A_r'']$  by removing row  $x_2$ , multiplying column  $y_1$  by  $-1$ , and swapping the order of columns  $y_0$  and  $y_1$ . Since  $B_r''$  is TU and since TUness is preserved under pivoting, taking submatrices, multiplying columns by  $\pm 1$  factors, and re-ordering columns, we conclude that  $[c_1'' \ c_2'' \ A_r'']$  is TU.

- (5) Let  $V$  be a square submatrix of  $[c_0'' \ c_1'' \ c_2'' \ A_r'']$ . Our goal is to show that  $\det V \in \{0, \pm 1\}$ .

Suppose that column  $c_2''$  is not in  $V$ . Then  $V$  is a submatrix of  $B_r''$ , which is TU. Thus,  $\det V \in \{0, \pm 1\}$ . Going forward we assume that column  $z$  is in  $V$ .

Suppose that columns  $c_0''$  and  $c_1''$  are both in  $V$ . Then  $V$  contains columns  $c_0''$ ,  $c_1''$ , and  $c_2'' = c_0'' - c_1''$ , which are linearly. Thus,  $\det V = 0$ . Going forward we assume that at least one of the columns  $c_0''$  and  $c_1''$  is not in  $V$ .

Suppose that column  $c_1''$  is not in  $V$ . Then  $V$  is a submatrix of  $[c_0'' \ c_2'' \ A_r'']$ , which is TU by item 3. Thus,  $\det V \in \{0, \pm 1\}$ . Similarly, if column  $c_0''$  is not in  $V$ , then  $V$  is a submatrix of  $[c_1'' \ c_2'' \ A_r'']$ , which is TU by item 4. Thus,  $\det V \in \{0, \pm 1\}$ .

□

**Lemma 56.** Suppose that  $B_\ell$  from Definition 46 has a TU signing  $B_\ell'$ . Let  $B_\ell''$  be the canonical re-signing (from Definition 51) of  $B_\ell'$ . Let  $d_0'' = B_\ell''(x_0, Y_\ell)$ ,  $d_1'' = B_\ell''(x_1, Y_\ell)$ , and  $d_2'' = d_0'' - d_1''$ . Then the following statements hold.

- (1) For every  $j \in Y_\ell$ ,  $\begin{bmatrix} d_0''(j) \\ d_1''(j) \end{bmatrix} \in \{0, \pm 1\}^{\{x_1, x_0\}} \setminus \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$ .  
(2) For every  $j \in Y_\ell$ ,  $d_2''(j) \in \{0, \pm 1\}$ .  
(3)  $\begin{bmatrix} A_\ell'' \\ d_0'' \\ d_2'' \end{bmatrix}$  is TU.  
(4)  $\begin{bmatrix} A_\ell'' \\ d_1'' \\ d_2'' \end{bmatrix}$  is TU.

$$(5) \begin{bmatrix} A''_\ell \\ d''_0 \\ d''_1 \\ d''_2 \end{bmatrix} \text{ is TU.}$$

*Proof.* Apply Lemma 55 to  $B_\ell^\top$ , or repeat the same arguments up to transposition.  $\square$

**Lemma 57.** Let  $B''$  be from Definition 53. Let  $c''_0 = B''(X_r, y_0)$ ,  $c''_1 = B''(X_r, y_1)$ , and  $c''_2 = c''_0 - c''_1$ . Similarly, let  $d''_0 = B''(x_0, Y_\ell)$ ,  $d''_1 = B''(x_1, Y_\ell)$ , and  $d''_2 = d''_0 - d''_1$ . Then the following statements hold.

- (1) For every  $i \in X_r$ ,  $c''_2(i) \in \{0, \pm 1\}$ .
- (2) If  $D''_0 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ , then  $D'' = c''_0 \otimes d''_0 - c''_1 \otimes d''_1$ . If  $D''_0 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ , then  $D'' = c''_0 \otimes d''_0 - c''_0 \otimes d''_1 + c''_1 \otimes d''_1$ .
- (3) For every  $j \in Y_\ell$ ,  $D''(X_r, j) \in \{0, \pm c''_0, \pm c''_1, \pm c''_2\}$ .
- (4) For every  $i \in X_r$ ,  $D''(i, Y_\ell) \in \{0, \pm d''_0, \pm d''_1, \pm d''_2\}$ .
- (5)  $\begin{bmatrix} A''_\ell \\ D'' \end{bmatrix}$  is TU.

*Proof.*

- (1) Holds by Lemma 55.2.
- (2) Note that

$$\begin{bmatrix} D''_\ell \\ D''_{\ell r} \end{bmatrix} = \begin{bmatrix} D''_0 \\ D''_r \end{bmatrix} \cdot (D''_0)^{-1} \cdot D''_\ell, \quad \begin{bmatrix} D''_0 \\ D''_r \end{bmatrix} = \begin{bmatrix} D''_0 \\ D''_r \end{bmatrix} \cdot (D''_0)^{-1} \cdot D''_0, \quad \begin{bmatrix} D''_0 \\ D''_r \end{bmatrix} = \begin{bmatrix} c''_0 & c''_1 \end{bmatrix}, \quad \begin{bmatrix} D''_\ell & D''_0 \end{bmatrix} = \begin{bmatrix} d''_0 \\ d''_1 \end{bmatrix}.$$

Thus,

$$D'' = \begin{bmatrix} D''_\ell & D''_0 \\ D''_{\ell r} & D''_r \end{bmatrix} = \begin{bmatrix} D''_0 \\ D''_r \end{bmatrix} \cdot (D''_0)^{-1} \cdot \begin{bmatrix} D''_\ell & D''_0 \end{bmatrix} = \begin{bmatrix} c''_0 & c''_1 \end{bmatrix} \cdot (D''_0)^{-1} \cdot \begin{bmatrix} d''_0 \\ d''_1 \end{bmatrix}.$$

Considering the two cases for  $D''_0$  and performing the calculations yields the desired results.

- (3) Let  $j \in Y_\ell$ . By Lemma 56.1,  $\begin{bmatrix} d''_0(j) \\ d''_1(j) \end{bmatrix} \in \{0, \pm 1\}^{\{x_1, x_0\}} \setminus \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$ . Consider two cases.
  - (a) If  $D''_0 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ , then by item 2 we have  $D''(X_r, j) = d''_0(j) \cdot c''_0 + (-d''_1(j)) \cdot c''_1$ .  
By considering all possible cases for  $d''_0(j)$  and  $d''_1(j)$ , we conclude that  $D''(X_r, j) \in \{0, \pm c''_0, \pm c''_1, \pm(c''_0 - c''_1)\}$ .
  - (b) If  $D''_0 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ , then by item 2 we have  $D''(X_r, j) = (d''_0(j) - d''_1(j)) \cdot c''_0 + d''_1(j) \cdot c''_1$ .  
By considering all possible cases for  $d''_0(j)$  and  $d''_1(j)$ , we conclude that  $D''(X_r, j) \in \{0, \pm c''_0, \pm c''_1, \pm(c''_0 - c''_1)\}$ .
- (4) Let  $i \in X_r$ . By Lemma 55.1,  $\begin{bmatrix} c''_0(i) & c''_1(i) \end{bmatrix} \in \{0, \pm 1\}^{\{y_0, y_1\}} \setminus \{[1 \ -1], [-1 \ 1]\}$ . Consider two cases.
  - (a) If  $D''_0 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ , then by item 2 we have  $D''(i, Y_\ell) = c''_0(i) \cdot d''_0 + (-c''_1(i)) \cdot d''_1$ .  
By considering all possible cases for  $c''_0(i)$  and  $c''_1(i)$ , we conclude that  $D''(i, Y_\ell) \in \{0, \pm d''_0, \pm d''_1, \pm d''_2\}$ .
  - (b) If  $D''_0 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ , then by item 2 we have  $D''(i, Y_\ell) = c''_0(i) \cdot d''_0 + (c''_1(i) - c''_0(i)) \cdot d''_1$ .  
By considering all possible cases for  $c''_0(i)$  and  $c''_1(i)$ , we conclude that  $D''(i, Y_\ell) \in \{0, \pm d''_0, \pm d''_1, \pm d''_2\}$ .

(5) By Lemma 56.5,  $\begin{bmatrix} A''_\ell \\ d''_0 \\ d''_1 \\ d''_2 \end{bmatrix}$  is TU. Since TUness is preserved under adjoining zero rows,

copies of existing rows, and multiplying rows by  $\pm 1$  factors,  $\begin{bmatrix} A''_\ell \\ 0 \\ \pm d''_0 \\ \pm d''_1 \\ \pm d''_2 \end{bmatrix}$  is also TU. By

item 4,  $\begin{bmatrix} A''_\ell \\ D'' \end{bmatrix}$  is a submatrix of the latter matrix, hence it is also TU.

□

#### 5.4. Proof of Regularity.

**Definition 58.** Let  $X'_\ell, Y'_\ell, X'_r, Y'_r$  be sets and let  $x_0$  and  $x_1$  be distinct elements contained neither in  $X'_\ell$  nor  $X'_r$ . Additionally, let  $c_0, c_1 \in \mathbb{Q}^{X'_r \cup \{x_1, x_0\}}$  be column vectors. We define

$\mathcal{C}(X'_\ell, Y'_\ell, X'_r, Y'_r; c_0, c_1)$  to be the family of matrices of the form  $\begin{bmatrix} A_\ell & 0 \\ D & A_r \end{bmatrix}$  such that:

- (1)  $A_\ell \in \mathbb{Q}^{X'_\ell \times Y'_\ell}$ ,  $A_r \in \mathbb{Q}^{(X'_r \cup \{x_1, x_0\}) \times Y'_r}$ , and  $D \in \mathbb{Q}^{(X'_r \cup \{x_1, x_0\}) \times Y'_\ell}$ ;
- (2)  $\begin{bmatrix} A_\ell \\ D \end{bmatrix}$  is TU;
- (3) for every  $j \in Y'_\ell$ ,  $D(X'_r, j) \in \{0, \pm c_0, \pm c_1, \pm(c_0 - c_1)\}$ ;
- (4)  $\begin{bmatrix} c_0 & c_1 & c_0 - c_1 & A_r \end{bmatrix}$  is TU;
- (5)  $\begin{bmatrix} A_\ell & 0 \\ D(x_0, Y'_\ell) & 1 \\ D(x_1, Y'_\ell) & 1 \end{bmatrix}$  is TU;
- (6)  $c_0(x_0) = 1$  and  $c_0(x_1) = 0$ ;
- (7) either  $c_1(x_0) = 0$  and  $c_1(x_1) = -1$ , or  $c_1(x_0) = 1$  and  $c_1(x_1) = 1$ .

**Lemma 59.** Let  $B''$  be from Definition 53. Then  $B'' \in \mathcal{C}(X'_\ell, Y'_\ell, X'_r, Y'_r; c''_0, c''_1)$  with  $X'_\ell = X_\ell \setminus \{x_1, x_0\}$ ,  $X'_r = X_r \setminus \{x_2, x_1, x_0\}$ ,  $Y'_\ell = Y_\ell \setminus \{y_2\}$ ,  $Y'_r = Y_r \setminus \{y_0, y_1\}$ ,  $x_0$  and  $x_1$  are the same,  $c''_0 = B''(X'_r, y_0)$ , and  $c''_1 = B''(X'_r, y_1)$ .

*Proof.* Recall that  $c''_0 - c''_1 \in \{0, \pm 1\}^{X'_r}$  by Lemma 57.1, so  $\mathcal{C}(X'_\ell, Y'_\ell, X'_r, Y'_r; c''_0, c''_1)$  is well-defined. To see that  $B'' \in \mathcal{C}(X'_\ell, Y'_\ell, X'_r, Y'_r; c''_0, c''_1)$ , note that all properties from Definition 58 are satisfied: property 3 holds by Lemma 57.3, property 4 holds by Lemma 55.5, and property 2 holds by Lemma 57.5. □

**Lemma 60.** Let  $C \in \mathcal{C}(X'_\ell, Y'_\ell, X'_r, Y'_r; c_0, c_1)$  from Definition 58. Let  $x \in X'_\ell$  and  $y \in Y'_\ell$  be such that  $A_\ell(x, y) \neq 0$ , and let  $C'$  be the result of performing a short tableau pivot in  $C$  on  $(x, y)$ . Then  $C' \in \mathcal{C}(X'_\ell, Y'_\ell, X'_r, Y'_r; c_0, c_1)$ .

*Proof.* Our goal is to show that  $C'$  satisfies all properties from Definition 58. Let  $C' = \begin{bmatrix} C'_{11} & C'_{12} \\ C'_{21} & C'_{22} \end{bmatrix}$ , and let  $\begin{bmatrix} A'_\ell \\ D' \end{bmatrix}$  be the result of performing a short tableau pivot on  $(x, y)$  in  $\begin{bmatrix} A_\ell \\ D \end{bmatrix}$ . Observe the following.

- By Lemma 13,  $C'_{11} = A'_\ell$ ,  $C'_{12} = 0$ ,  $C'_{21} = D'$ , and  $C'_{22} = A_r$ .
- Since  $\begin{bmatrix} A_\ell \\ D \end{bmatrix}$  is TU by property 2 for  $C$ , all entries of  $A_\ell$  are in  $\{0, \pm 1\}$ .
- $A_\ell(x, y) \in \{\pm 1\}$ , as  $A_\ell(x, y) \in \{0, \pm 1\}$  by the above observation and  $A_\ell(x, y) \neq 0$  by the assumption.
- Since  $\begin{bmatrix} A_\ell \\ D \end{bmatrix}$  is TU by property 2 for  $C$ , and since pivoting preserves TUness,  $\begin{bmatrix} A'_\ell \\ D' \end{bmatrix}$  is also TU.

These observations immediately imply properties 4 and 2 for  $C'$ . Indeed, property 4 holds for  $C'$ , since  $C'_{22} = A_r$  and  $\begin{bmatrix} c_0 & c_1 & c_0 - c_1 & A_r \end{bmatrix}$  is TU by property 4 for  $C$ . On the other hand, property 2 follows from  $C'_{11} = A'_\ell$ ,  $C'_{21} = D'$ , and  $\begin{bmatrix} A'_\ell \\ D' \end{bmatrix}$  being TU. Thus, it only remains to show that  $C'$  satisfies property 3. Let  $j \in Y_r$ . Our goal is to prove that  $D'(X'_r, j) \in \{0, \pm c_0, \pm c_1, \pm(c_0 - c_1)\}$ .

Suppose  $j = y$ . By the pivot formula,  $D'(X'_r, y) = -\frac{D(X'_r, y)}{A_\ell(x, y)}$ . Since  $D(X'_r, y) \in \{0, \pm c_0, \pm c_1, \pm(c_0 - c_1)\}$  by property 3 for  $C$  and since  $A_\ell(x, y) \in \{\pm 1\}$ , we get  $D'(X'_r, y) \in \{0, \pm c_0, \pm c_1, \pm(c_0 - c_1)\}$ .

Now suppose  $j \in Y_\ell \setminus \{y\}$ . By the pivot formula,  $D'(X'_r, j) = D(X'_r, j) - \frac{A_\ell(x, j)}{A_\ell(x, y)} \cdot D(X'_r, y)$ . Here  $D(X'_r, j), D(X'_r, y) \in \{0, \pm c_0, \pm c_1, \pm(c_0 - c_1)\}$  by property 3 for  $C$ , and  $A_\ell(x, j) \in \{0, \pm 1\}$  and  $A_\ell(x, y) \in \{\pm 1\}$  by the prior observations. Perform an exhaustive case distinction on  $D(X'_r, j), D(X'_r, y), A_\ell(x, j)$ , and  $A_\ell(x, y)$ . The number of cases can be significantly reduced by using symmetries. In every remaining case, we can either show that  $D'(X'_r, j) \in \{0, \pm c_0, \pm c_1, \pm(c_0 - c_1)\}$ , as desired, or obtain a contradiction with property 5.  $\square$

**Lemma 61.** Let  $C \in \mathcal{C}(X'_\ell, Y'_\ell, X'_r, Y'_r; c_0, c_1)$  from Definition 58. Then  $C$  is TU.

*Proof.* By Lemma 7, it suffices to show that  $C$  is  $k$ -PU for every  $k \in \mathbb{N}$ . We prove this claim by induction on  $k$ . The base case with  $k = 1$  holds, since properties 4 and 2 in Definition 58 imply that  $A_\ell, A_r$ , and  $D$  are TU, so all their entries of  $C = \begin{bmatrix} A_\ell & 0 \\ D & A_r \end{bmatrix}$  are in  $\{0, \pm 1\}$ , as desired.

Suppose that for some  $k \in \mathbb{N}$  we know that every  $C' \in \mathcal{C}(X'_\ell, Y'_\ell, X'_r, Y'_r; c_0, c_1)$  is  $k$ -PU. Our goal is to show that  $C$  is  $(k + 1)$ -PU, i.e., that every  $(k + 1) \times (k + 1)$  submatrix  $S$  of  $C$  has  $\det V \in \{0, \pm 1\}$ .

First, suppose that  $V$  has no rows in  $X'_\ell$ . Then  $V$  is a submatrix of  $\begin{bmatrix} D & A_r \end{bmatrix}$ , which is TU by property 4 in Definition 58, so  $\det V \in \{0, \pm 1\}$ . Thus, we may assume that  $S$  contains a row  $x_\ell \in X'_\ell$ .

Next, note that without loss of generality we may assume that there exists  $y_\ell \in Y'_\ell$  such that  $V(x_\ell, y_\ell) \neq 0$ . Indeed, if  $V(x_\ell, y) = 0$  for all  $y$ , then  $\det V = 0$  and we are done, and  $V(x_\ell, y) = 0$  holds whenever  $y \in Y'_r$ .

Since  $C$  is 1-PU, all entries of  $V$  are in  $\{0, \pm 1\}$ , and hence  $V(x_\ell, y_\ell) \in \{\pm 1\}$ . Thus, by Lemma 14, performing a short tableau pivot in  $V$  on  $(x_\ell, y_\ell)$  yields a matrix that contains a  $k \times k$  submatrix  $S''$  such that  $|\det V| = |\det S''|$ . Since  $V$  is a submatrix of  $C$ , matrix  $S''$  is a submatrix of the matrix  $C'$  resulting from performing a short tableau pivot in  $C$  on the same entry  $(x_\ell, y_\ell)$ . By Lemma 60, we have  $C' \in \mathcal{C}(X'_\ell, Y'_\ell, X'_r, Y'_r; c_0, c_1)$ . Thus, by the inductive hypothesis applied to  $S''$  and  $C'$ , we have  $\det S'' \in \{0, \pm 1\}$ . Since  $|\det V| = |\det S''|$ , we conclude that  $\det V \in \{0, \pm 1\}$ .  $\square$

**Lemma 62.**  $B''$  from Definition 53 is TU.

*Proof.* Combine the results of Lemmas 59 and 61.  $\square$

**Theorem 63.** Let  $M$  be a 3-sum of regular matroids  $M_\ell$  and  $M_r$ . Then  $M$  is also regular.

*Proof.* Let  $B_\ell, B_r$ , and  $B$  be standard  $\mathbb{Z}_2$  representation matrices from Definition 47. Since  $M_\ell$  and  $M_r$  are regular, by Lemma 34,  $B_\ell$  and  $B_r$  have TU signings. Then the canonical signing  $B''$  from Definition 53 is a TU signing of  $B$ . Indeed,  $B''$  is a signing of  $B$  by Lemma 54, and  $B''$  is TU by Lemma 62. Thus,  $M$  is regular by Lemma 34.  $\square$

## 6. SPECIAL MATROIDS

**Definition 64.** Let  $A \in \mathbb{Q}^{X \times Y}$  be a matrix. If for all  $j \in Y$ , one has that  $a_{i,j} = 0$  for all  $i \in X$ , or that there exists  $i_1, i_2 \in X$  such that

$$a_{i,j} = \begin{cases} 1 & \text{if } i = i_1 \\ -1 & \text{if } i = i_2 \\ 0 & \text{otherwise,} \end{cases}$$

then we call  $A$  a node-incidence matrix for a (directed) graph whose nodes are indexed by  $X$  and whose edges are indexed by  $Y$ .

**Definition 65.** We say that a matroid is graphic if it can be represented by a node-incidence matrix.

**Definition 66.** Let  $S$  be a standard representation given by matrix  $B$ . The dual of  $S$  is given by  $-B^\top$ .

**Definition 67.** We say a matroid is co-graphic if its dual is graphic.

**Definition 68.** The matroid with standard representation

$$\begin{bmatrix} 1 & 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \end{bmatrix}$$

over  $\mathbb{Z}_2$  is called  $R_{10}$ .

**Theorem 69.** *The matroid  $R_{10}$  is regular.*

*Proof.* See Lean implementation. □

**Theorem 70.** *Every graphic matroid is regular.*

*Proof.* See Lean implementation. □

### 6.1. Regularity of Cographic Matroids.

**Lemma 71** (Row space of a standard representation). Let  $X$  and  $Y$  be disjoint finite sets and let

$$B \in \mathbb{F}_2^{X \times Y}.$$

Consider the matrix

$$A := [\mathbf{1}_x \mid B] \in \mathbb{F}_2^{X \times (X \cup Y)},$$

where the columns are indexed by  $E := X \cup Y$  and the rows by  $X$ . Then the row space of  $A$  is

$$\text{row}(A) = \{(u, uB) \mid u \in \mathbb{F}_2^X\} \subseteq \mathbb{F}_2^X \oplus \mathbb{F}_2^Y \cong \mathbb{F}_2^E.$$

*Proof.* The  $x$ -th row of  $A$  is  $(e_x, B_{x,*})$ , where  $e_x$  is the standard basis vector in  $\mathbb{F}_2^X$  and  $B_{x,*}$  is the  $x$ -th row of  $B$ . A general linear combination of the rows is therefore

$$\sum_{x \in X} u_x (e_x, B_{x,*}) = \left(u, \sum_{x \in X} u_x B_{x,*}\right) = (u, uB),$$

where  $u = (u_x)_{x \in X} \in \mathbb{F}_2^X$ . Conversely, every pair  $(u, uB)$  arises in this way, so these are exactly the row vectors. □

**Lemma 72** (Orthogonal complement of a standard row space). Let  $A = [\mathbf{1}_x \mid B]$  be as in Lemma 71, and let

$$U := \text{row}(A) \subseteq \mathbb{F}_2^{X \cup Y}.$$

Then the orthogonal complement of  $U$  is

$$U^\perp = \{(bB^\top, b) \mid b \in \mathbb{F}_2^Y\}.$$

Equivalently, if  $B^* := -B^\top$ , then

$$U^\perp = \{(bB^*, b) \mid b \in \mathbb{F}_2^Y\}.$$

*Proof.* Write vectors in  $\mathbb{F}_2^{X \cup Y}$  as pairs  $(a, b)$  with  $a \in \mathbb{F}_2^X$  and  $b \in \mathbb{F}_2^Y$ . By Lemma 71, any element of  $U$  has the form  $(u, uB)$  with  $u \in \mathbb{F}_2^X$ . The orthogonality condition  $(a, b) \in U^\perp$  means

$$\begin{aligned} 0 &= (a, b) \cdot (u, uB) \\ &= a \cdot u + b \cdot (uB) \\ &= a \cdot u + (bB^\top) \cdot u \\ &= (a + bB^\top) \cdot u \end{aligned}$$

for all  $u \in \mathbb{F}_2^X$ . Hence we must have  $a = bB^\top$ , and then

$$U^\perp = \{ (bB^\top, b) \mid b \in \mathbb{F}_2^Y \}.$$

Over  $\mathbb{F}_2$  we have  $-1 = 1$ , so  $B^* = -B^\top = B^\top$ , yielding the alternative description.  $\square$

**Lemma 73** (Row space of the dual standard matrix). With  $B$  and  $B^* = -B^\top$  as above, define

$$A^* := [\mathbf{1}_Y \mid B^*] \in \mathbb{F}_2^{Y \times (X \cup Y)}.$$

Then

$$\text{row}(A^*) = U^\perp,$$

where  $U = \text{row}(A)$  and  $U^\perp$  is given by Lemma 72.

*Proof.* The  $y$ -th row of  $A^*$  is  $(e_y, B_{y,*}^*)$  with  $e_y \in \mathbb{F}_2^Y$ . A general linear combination of the rows is

$$\sum_{y \in Y} b_y (e_y, B_{y,*}^*) = (b, bB^*),$$

where  $b = (b_y)_{y \in Y} \in \mathbb{F}_2^Y$ . Thus

$$\text{row}(A^*) = \{ (b, bB^*) \mid b \in \mathbb{F}_2^Y \}.$$

Identifying  $\mathbb{F}_2^{X \cup Y}$  as  $\mathbb{F}_2^X \oplus \mathbb{F}_2^Y$  with coordinates ordered as  $(X, Y)$ , this is exactly the set

$$\{ (bB^*, b) \mid b \in \mathbb{F}_2^Y \},$$

which coincides with  $U^\perp$  by Lemma 72.  $\square$

**Lemma 74** (Dual vector matroid via orthogonal complement). Let  $A$  and  $A'$  be matrices over a field  $F$  with the same column index set  $E$ , and suppose

$$\text{row}(A') = \text{row}(A)^\perp \subseteq F^E.$$

Let  $M(A)$  and  $M(A')$  be the vector matroids represented by  $A$  and  $A'$ . Then

$$M(A') = M(A)^*.$$

*Proof.* Let  $F \subseteq E$ .

( $\Rightarrow$ ) Suppose  $F$  is dependent in  $M(A)$ . Then there exists a nonzero vector  $c \in F^F$  such that  $A_F c = 0$ . Extend  $c$  by zero outside  $F$  (still denoted  $c$ ). The condition  $Ac = 0$  means each row  $r$  of  $A$  satisfies  $r \cdot c = 0$ , hence  $c \in \text{row}(A)^\perp = \text{row}(A')$ . Write

$$c = \sum_i \lambda_i r'_i,$$

where the  $r'_i$  are rows of  $A'$  and not all  $\lambda_i$  are zero. For every  $e \in E \setminus F$  we have  $c_e = 0$ , so

$$\left( \sum_i \lambda_i r'_i \right) \Big|_{E \setminus F} = 0.$$

Hence the rows of  $A'$  indexed by  $E \setminus F$  admit a nontrivial linear combination giving the zero row, so  $E \setminus F$  is dependent in  $M(A')$ .

( $\Leftarrow$ ) The same argument with  $A$  and  $A'$  interchanged, using  $\text{row}(A) = (\text{row}(A')^\perp)$ , shows that if  $E \setminus F$  is dependent in  $M(A')$ , then  $F$  is dependent in  $M(A)$ .

Thus

$$F \text{ dependent in } M(A) \iff E \setminus F \text{ dependent in } M(A'),$$

which is the defining property of duality.  $\square$

**Theorem 75** (Dual of standard representation corresponds to dual matroid). *Let  $M$  be a binary matroid on ground set  $E = X \cup Y$ , with standard representation  $B$  so that*

$$A = [\mathbf{1}_X \mid B].$$

Let  $B^* := -B^\top$  and

$$A^* := [\mathbf{1}_Y \mid B^*].$$

Then  $M(A^*) = M(A)^* = M^*$ .

*Proof.* By Lemma 71 and Lemma 72, if  $U = \text{row}(A)$  then  $U^\perp$  has the form

$$U^\perp = \{ (bB^*, b) \mid b \in \mathbb{F}_2^Y \}.$$

By Lemma 73, we have

$$\text{row}(A^*) = U^\perp = \text{row}(A)^\perp.$$

Therefore, by Lemma 74, the column-matroid  $M(A^*)$  is the dual of  $M(A)$ :

$$M(A^*) = M(A)^* = M^*.$$

$\square$

**Lemma 76.** The dual matroid of a regular matroid is also a regular matroid.

*Proof.* Let  $M$  be a regular matroid. We wish to show that  $M^*$  is also regular.

Take a standard  $\mathbb{Z}_2$ -representation matrix  $B$  of  $M$ . By Lemma 34, since  $M$  is regular, there exists a TU signing  $B'$  of  $B$ :  $B'$  is a matrix over  $\mathbb{Q}$  that is TU, and  $|B'(i, j)| = B(i, j)$  for all entries. So  $M$  is represented (over  $\mathbb{Q}$ ) by a TU matrix  $B'$  whose pattern of zero and non-zero entries is exactly that of  $B$ .

From Theorem 75, if a matroid  $M$  has standard representation matrix  $B$ , then its dual  $M^*$  has the standard representation matrix  $B^* = -B^\top$ . The TU signing of this dual standard matrix,  $(B')^* = -(B')^\top$ , preserves total unimodularity, so  $(B')^*$  is a TU matrix whose support is exactly  $B^*$ .

Since we have just exhibited a TU signing of  $M^*$  (i.e.,  $(B')^*$ ), the dual matroid  $M^*$  is regular by Lemma 34.  $\square$

**Theorem 77.** Every cographic matroid is regular.

*Proof.* We know that all graphic matroids are regular by Theorem 70. Recall that we say a matroid is cographic if its dual is graphic. So it suffices to show regularity is preserved under duals, which we showed in Lemma 76.  $\square$

## 7. CONCLUSION

**Definition 78.** Any graphic matroid is good. Any cographic matroid is good. Any matroid isomorphic to R10 is good. Any 1-sum (in the sense of Definition 36) of good matroids is a good matroid. Any 2-sum (in the sense of Definition 41) of good matroids is a good matroid. Any 3-sum (in the sense of Definition 47) of good matroids is a good matroid.

**Corollary 79.** Any good matroid is regular. This is a corollary of the easy direction of the Seymour theorem.

*Proof.* Structural induction.  $\square$

## REFERENCES

- [1] K. Truemper, *Matroid Decomposition*, Leibniz Center for Informatics, 2016. Available at <https://www2.math.ethz.ch/EMIS/monographs/md/>.
- [2] J. Oxley, *Matroid Theory*, 2nd ed., Oxford University Press, 2011. <https://doi.org/10.1093/acprof:oso/9780198566946.001.0001>.
- [3] H. Bruhn, R. Diestel, M. Kriesell, R. Pendavingh, and P. Wollan, *Axioms for Infinite Matroids*, arXiv preprint, 2013. Available at <https://arxiv.org/abs/1303.5277>.

IVAN SERGEEV, INSTITUTE OF SCIENCE AND TECHNOLOGY AUSTRIA  
*Email address:* `ivan.sergeev@ist.ac.at`

MARTIN DVORAK, INSTITUTE OF SCIENCE AND TECHNOLOGY AUSTRIA  
*Email address:* `martin.dvorak@ista.ac.at`

MARK SANDEY  
*Email address:* `mark@san dey-family.com`

PIETRO MONTICONE, UNIVERSITY OF TRENTO

CAMERON RAMPELL  
*Email address:* `cameronrampell@gmail.com`