Formalizing Brownian motion

April 27, 2024

Our goal is to write down the steps necessary in order to formalize Brownian motions (or \mathbb{R}^d -valued Gaussian processes) in some generality using mathlib.

Remark 0.1 (Notation). We will write (E, r) for some extended pseudo-metric space, $\mathcal{P}(E)$ for the set of probability measures on the Borel σ -algebra on E, $\Bbbk \in \{\mathbb{R}, \mathbb{C}\}$, and $\mathcal{C}_b(E, \Bbbk)$ the set of \Bbbk -valued bounded continuous functions on E. For some $\mathbf{P} \in \mathcal{P}(E)$ and $f \in \mathcal{C}_b(E, \Bbbk)$, we let $\mathbf{P}[f] := \int f(x)\mathbf{P}(dx) \in \Bbbk$ be the expectation.

0 Some simple probability results

The following is a simple consequence of dominated convergence, and often needed in probability theory.

Definition 0.1. Let E be some set and $f, f_1, f_2, \ldots : E \to \Bbbk$. We say that $f_1, f_2, \ldots :$ converges to f boundedly pointwise if $f_n \xrightarrow{n \to \infty} f$ pointwise and $\sup_n ||f_n|| < \infty$. We write $f_n \xrightarrow{n \to \infty} b_p f$

Proof. Note that the constant function $x \mapsto \sup_n ||f_n||$ is integrable (since **P** is finite), so the result follows from dominated convergence.

Definition 0.3. Let $X, X_1, X_2, ..., all E$ -valued random variables.

- 1. We say that $X_n \xrightarrow{n \to \infty} X$ almost everywhere if $\mathbf{P}(\lim_{n \to \infty} X_n = X) = 1$. We also write $X_n \xrightarrow{n \to \infty}_{ae} X$.
- 2. We say that $X_n \xrightarrow{n \to \infty} X$ in probability (or in measure) if, for all $\varepsilon > 0$,

$$\lim_{n \to \infty} \mathbf{P}(r(X_n, X) > \varepsilon) = 0.$$

The two notions here are denoted $\forall^m (x : \alpha) \partial P$, Filter.Tendsto (fun n => X n x) Filter.atTop (nhds (X x)) and MeasureTheory.TendstoInMeasure, respectively.

Lemma 0.4. Let $X, X_1, X_2, ...$ be E-valued random variables with $X_n \xrightarrow{n \to \infty}_{ae} X$. Then, $X_n \xrightarrow{n \to \infty}_p X$.

This result is called MeasureTheory.tendstoInMeasure_of_tendsto_ae in mathlib. We also need the (almost sure) uniquess of the limit in measure, which is not formalized in mathlib yet:

Lemma 0.5 (Uniqueness of a limit in probability). Let $X, Y, X_1, X_2, ...$ be *E*-valued random variables with $X_n \xrightarrow{n \to \infty}_p X$ and $X_n \xrightarrow{n \to \infty}_p Y$. Then, X = Y, almost surely.

Proof. We write, using monotone convergence and Lemma ??

$$\mathbf{P}(X \neq Y) = \lim_{\varepsilon \downarrow 0} \mathbf{P}(r(X, Y) > \varepsilon) \le \lim_{\varepsilon \downarrow 0} \lim_{n \to \infty} \mathbf{P}(r(X, X_n) > \varepsilon/2) + \mathbf{P}(r(Y, X_n) > \varepsilon/2) = 0.$$

Lemma 0.6. Let I be some (finite or infinite) set and $(X_t)_{t \in I}$ be a family of random variables with values in $[0, \infty)$. Then, $\sup_{t \in I} X_t \leq \sum_{t \in I} X_t$.

1 Separating algebras and characteristic functions

Definition 1.1 (Separating class of functions). Let $\mathcal{M} \subseteq \mathcal{C}_b(E, \mathbb{k})$.

- 1. If, for all $x, y \in E$ with $x \neq y$, there is $f \in \mathcal{M}$ with $f(x) \neq f(y)$, we say that \mathcal{M} separates points.
- 2. If, for all $\mathbf{P}, \mathbf{Q} \in \mathcal{P}(E)$,

$$\mathbf{P} = \mathbf{Q} \quad iff \quad \mathbf{P}[f] = \mathbf{Q}[f] \text{ for all } f \in \mathcal{M},$$

we say that \mathcal{M} is separating in $\mathcal{P}(E)$.

3. If (i) $1 \in \mathcal{M}$ and (ii) if \mathcal{M} is closed under sums and products, then we call \mathcal{M} a (sub-)algebra. If $\Bbbk = \mathbb{C}$, and (iii) if \mathcal{M} is closed under complex conjugation, we call \mathcal{M} a star-(sub-)algebra.

In mathlib, 1. and 3. of the above definition are already implemented:

```
structure Subalgebra (R : Type u) (A : Type v) [CommSemiring R] [Semiring A]
[Algebra R A] extends Subsemiring :
Type v
```

```
abbrev Subalgebra.SeparatesPoints {\alpha : Type u_1} [TopologicalSpace \alpha]
{R : Type u_2} [CommSemiring R] {A : Type u_3} [TopologicalSpace A]
[Semiring A] [Algebra R A] [TopologicalSemiring A] (s : Subalgebra R C(\alpha, A))
: Prop
```

The latter is an extension of Set.SeparatesPoints, which works on any set of functions. For the first result, we already need that (E, r) has a metric structure. There is a formalization of this result in https://github.com/pfaffelh/some_probability/tree/master.

I:unique Lemma 1.2. $\mathcal{M} := \mathcal{C}_b(E, \mathbb{k})$ is separating.

Proof. We restrict ourselves to $\mathbf{k} = \mathbb{R}$, since the result for $\mathbf{k} = \mathbb{C}$ follows by only using functions with vanishing imaginary part. Let $\mathbf{P}, \mathbf{Q} \in \mathcal{P}(E)$. We will prove that $\mathbf{P}(A) = \mathbf{Q}(A)$ for all A closed. Since the set of closed sets is a π -system generating the Borel- σ -algebra, this suffices for $\mathbf{P} = \mathbf{Q}$. So, let A be closed and $g = 1_A$ be the indicator function. Let $g_n(x) := (1 - nr(A, x))^+$ (where $r(A, y) := \inf_{y \in A} r(y, x)$) and note that $g_n(x) \xrightarrow{n \to \infty} 1_A(x)$. Then, we have by dominated convergence

$$\mathbf{P}(A) = \lim_{n \to \infty} \mathbf{P}[g_n] = \lim_{n \to \infty} \mathbf{Q}[g_n] = \mathbf{Q}(A),$$

and we are done.

We will use the Stone-Weierstrass Theorem below. Here is its version in mathlib. Note that this requires E to be compact.

theorem ContinuousMap.starSubalgebra_topologicalClosure_eq_top_of_separatesPoints {k : **Type** u_2} {X : **Type** u_1} [IsROrC k] [TopologicalSpace X] [CompactSpace X] (A : StarSubalgebra k C(X, k)) (hA : Subalgebra.SeparatesPoints A.toSubalgebra) : StarSubalgebra.topologicalClosure A = \top

We also need (as proved in the last project):

theorem innerRegular_isCompact_isClosed_measurableSet_of_complete_countable [PseudoEMetricSpace α] [CompleteSpace α] [SecondCountableTopology α] [BorelSpace α] (P : Measure α) [IsFiniteMeasure P] : P.InnerRegular (**fun** s => IsCompact s \wedge IsClosed s) MeasurableSet

The proof of the following result follows [?, Theorem 3.4.5].

T:wc3 Theorem 1 (Algebras separating points and separating algebras).

Let (E, r) be a complete and separable extended pseudo-metric space, and $\mathcal{M} \subseteq \mathcal{C}_b(E, \mathbb{k})$ be a star-sub-algebra that separates points. Then, \mathcal{M} is separating.

Proof. Let $\mathbf{P}, \mathbf{Q} \in \mathcal{P}(E)$, $\varepsilon > 0$ and K compact, such that $\mathbf{P}(K) > 1 - \varepsilon$, $\mathbf{Q}(K) > 1 - \varepsilon$, and $g \in \mathcal{C}_b(E, \mathbb{k})$. According to the Stone-Weierstrass Theorem, there is $(g_n)_{n=1,2,\dots}$ in \mathcal{M} with

$$\sup_{x \in K} |g_n(x) - g(x)| \xrightarrow{n \to \infty} 0. \tag{1} \quad \text{eq:wc9}$$

So, (note that $C := \sup_{x \ge 0} x e^{-x^2} < \infty$)

$$\begin{split} \left| \mathbf{P}[ge^{-\varepsilon g^2}] - \mathbf{Q}[ge^{-\varepsilon g^2}] \right| &\leq \left| \mathbf{P}[ge^{-\varepsilon g^2}] - \mathbf{P}[ge^{-\varepsilon g^2}; K] \right| \\ &+ \left| \mathbf{P}[ge^{-\varepsilon g^2}; K] - \mathbf{P}[g_n e^{-\varepsilon g^2_n}; K] \right| \\ &+ \left| \mathbf{P}[g_n e^{-\varepsilon g^2_n}; K] - \mathbf{P}[g_n e^{-\varepsilon g^2_n}] \right| \\ &+ \left| \mathbf{P}[g_n e^{-\varepsilon g^2_n}] - \mathbf{Q}[g_n e^{-\varepsilon g^2_n}] \right| \\ &+ \left| \mathbf{Q}[g_n e^{-\varepsilon g^2_n}] - \mathbf{Q}[g_n e^{-\varepsilon g^2_n}; K] \right| \\ &+ \left| \mathbf{Q}[g_n e^{-\varepsilon g^2_n}] - \mathbf{Q}[ge^{-\varepsilon g^2}; K] \right| \\ &+ \left| \mathbf{Q}[ge^{-\varepsilon g^2}; K] - \mathbf{Q}[ge^{-\varepsilon g^2}] \right| \end{split}$$

We bound the first term by

$$\left|\mathbf{P}[ge^{-\varepsilon g^2}] - \mathbf{P}[ge^{-\varepsilon g^2}; K]\right| \leq \frac{C}{\sqrt{\varepsilon}} \mathbf{P}(K^c) \leq C\sqrt{\varepsilon},$$

and analogously for the third, fifth and last. The second and second to last vanish for $n \to \infty$ due to (??). Since \mathcal{M} is an algebra, we can approximate, using dominated convergence,

$$\mathbf{P}[g_n e^{-\varepsilon g_n^2}] = \lim_{m \to \infty} \mathbf{P}[\underbrace{g_n \left(1 - \frac{\varepsilon g_n^2}{m}\right)^m}_{\in \mathcal{M}}] = \lim_{m \to \infty} \mathbf{Q}[\underbrace{g_n \left(1 - \frac{\varepsilon g_n^2}{m}\right)^m}_{\in \mathcal{M}}] = \mathbf{Q}[g_n e^{-\varepsilon g_n^2}],$$

so the fourth term vanishes for $n \to \infty$ as well. Concluding,

$$\left|\mathbf{P}[g] - \mathbf{Q}[g]\right| = \lim_{\varepsilon \to 0} \left|\mathbf{P}[ge^{-\varepsilon g^2}] - \mathbf{Q}[ge^{-\varepsilon g^2}]\right| \le 4C \lim_{\varepsilon \to 0} \sqrt{\varepsilon} = 0$$

Since g was arbitrary and $C_b(E, \mathbb{k})$ is separating by Lemma ??, we find $\mathbf{P} = \mathbf{Q}$.

We now come to characteristic functions and Laplace transforms.

Pr:charl | Proposition 1.3 (Charakteristic functions determine distributions uniquely).

A probability measure $\mathbf{P} \in \mathcal{P}(\mathbb{R}^d)$ is uniquely given by its characteristic function. In other words, if $\mathbf{P}, \mathbf{Q} \in \mathcal{P}(\mathbb{R}^d)$ are such that $\int e^{itx} \mathbf{P}(dx) = \int e^{itx} \mathbf{Q}(dx)$ for all $t \in \mathbb{R}^d$. Then, $\mathbf{P} = \mathbf{Q}$.

Proof. The set

$$\mathcal{M} := \left\{ x \mapsto \sum_{k=1}^{n} a_k e^{it_k x}; n \in \mathbb{N}, a_1, ..., a_n \in \mathcal{C}, t_1, ..., 1_n \in \mathbb{R}^d \right\}$$

separates points in \mathbb{R}^d . Since $\mathcal{M} \subseteq \mathcal{C}_b(\mathbb{R}^d, \mathbb{k})$ contains 1, is closed under sums and products, and closed under complex conjugation, it is a star-subalgebra of $\mathcal{C}_b(E, \mathbb{C})$. So, the assertion directly follows from Theorem ??.

rem:proj

Remark 1.4. We also need to show the following: For $J \subseteq I$, where I is finite, let ψ be the characteristic function for some distribution on \mathbb{R}^I . Then, for the projection π_J , the characteristic function of the image measure under π_J is given by $\psi \circ g_J$, where $(g_J(t)_j) = t_j$ for $j \in J$ and $(g(t)_j) = 0$ otherwise. In other words, when computing the characteristic function of a projection, just set the coordinates in $t \mapsto \psi(t)$ which need to be projected out to 0.

2 Gaussian random variables

Define an arbitrary family of Gaussian rvs with values in \mathbb{R}^d by (i) defining a standard normal distribution on \mathbb{R} with the correct density, (ii) show that its characteristic function is given by $\psi(t) = e^{-t^2/2}$, (iii) define an independent finite family of standard normal Gaussians using finite product measures and (iv) define a general independent family by taking some symmetric, positive definite $C \in \mathbb{R}^{d \times d}$, some¹ $A \in \mathbb{R}^{d \times d}$ with $C = A^{\top}A$, and define the Gaussian measure as the image measure of the independent family Y under the map $X = AY + \mu$. Show that

$$\mathbb{E}[e^{itX}] = \mathbb{E}[e^{it(\mu+AY)}] = e^{it\mu}\mathbb{E}[e^{itAY}] = e^{it\mu}\mathbb{E}\Big[\exp\left(i\sum_{kl}t_kA_{kl}Y_l\right)\Big]$$
$$= e^{it\mu}\prod_l \mathbb{E}\Big[\exp\left(i(\sum_k t_kA_{kl})Y_l\right)\Big] = e^{it\mu}\prod_l \mathbb{E}[e^{i(tA_{.l})Y_l}]$$
$$= e^{it\mu}\prod_l e^{-(tA_{.l})^2/2} = e^{it\mu}e^{-\sum_l(tA_{.l})(A_{l.}^{\top}t^{\top})/2} = e^{it\mu-tCt^{\top}/2}.$$

In particular, this shows that the distribution does not depend on the choice of A as long as $A^{\top}A = C$. Together with Proposition ??, this shows that there is a unique probability measure on \mathbb{R}^d with characteristic function $t \mapsto e^{it\mu - tCt^{\top}/2}$ for any vector μ and symmetric and positive definite matrix C.

3 Projectivity

S:proj

For projectivity of finite-dimensional distributions of the BM, proceed as follows: (i) For $I = \{s_1, ..., s_n\} \subseteq \mathbb{R}^d$ (with $s_1 < ... < s_d$), define P_J as the unique probability measure with characteristic function $\psi_I(t) = e^{-tC_I t^\top/2}$ with $C_{ij} = s_i \wedge s_j$. For $J \subseteq I$, we then have that the characteristic function of the projection to coordinates in J is (see Remark ??) $\psi_I \circ g_J = e^{-g_J(.)C_I g_J(.)^\top} = e^{-.C_J - /2} = \psi_J$. In other words, this is the required projectivity of $(P_I)_{I \subseteq f[0,\infty)}$.

4 The Kolmogorov-Chentsov criterion

In this section, let $(D, r_D), (E, r_E)$ be extended pseudo-metric spaces. In addition, we will only have a single probability measure in this section, so we write $\mathbf{P}(.)$ for probabilities and $\mathbf{E}[.]$ for its expectations.

Definition 4.1 (Local Hölder). Let $f: D \to E$ and $s \in D$. If there is $\tau > 0$ and some $C < \infty$ with $r_E(f(s), f(t)) \leq Cr_D(s, t)^{\gamma}$ for all t with $r_D(s, t) < \tau$, we call f locally Hölder of order γ at s.

Hölder is implemented as HolderOnWith (on a set) and HolderWith. Moreover, locally Hölder at a point is used for $\gamma = 1$ (i.e. Lipschitz continuity) e.g. in continuousAt_of_locally_lipschitz (Every function, which is locally Lipschitz at a point, is continuous.)

I:holderext Lemma 4.2. Let D, E be extended pseudo-metric spaces and $f: D \to E$ and $s \in D$.

- 1. If f is locally Hölder at x, it is continuous at x.
- If E is complete, A ⊆ D is dense, and g : A → E is Hölder, it can be extended to a Hölder-γ-function on D.

¹In order to see that such an A exists, consider some orthogonal O and a diagonal matrix D with $C = O^{\top}DO$ and set $A := \sqrt{D}O$, where \sqrt{D} is the diagonal matrix with entries $\sqrt{\lambda_i}$ for all eigenvalues λ_i of C. Then, $A^{\top}A = O^{\top}\sqrt{D}\sqrt{D}O = O^{\top}DO = C$.

Proof. 1. Since f is locally Hölder at s, choose $\tau > 0$ and $C < \infty$ such that $r_E(f(s), f(t)) \leq Cr(s,t)^{\gamma}$ for all t with $r_D(s,t) < \tau$. For $\varepsilon > 0$, there is $\delta' > 0$ such $r_D(s,t)^{\gamma} < \varepsilon/C$ for all $t \in B_{\delta'}(s)$. Choose $\delta := \tau \wedge \delta'$ in order to see, for $t \in B_{\delta}(s)$

$$r_E(f(s), f(t)) \le Cr(s, t)^{\gamma} < \varepsilon$$

2. For $s \in D$, choose $s_1, s_2, ... \in A$ with $s_n \xrightarrow{n \to \infty} s$. Then, note that $r_E(f(s_n), f(s_M)) \leq Cr_D(s_n, s_m) \xrightarrow{m, n \to \infty} 0$, so $(f(s_n))_{n=1,2,...}$ is a Cauchy-sequence in E. We define f(s) to be its limit. Then, for $s, t \in D$ and the sequences $s_1, s_2, ... \in D, t_1, t_2, ... \in D$ with $s_n \xrightarrow{n \to \infty} s, t_n \xrightarrow{n \to \infty} t$,

$$r_E(f(s), f(t)) = \lim_{n \to \infty} r_E(f(s_n), f(t_n)) \le \lim_{n \to \infty} Cr_D(s_n, t_n) = Cr_D(s, t).$$

For 1., continuous At_of_locally_lipschitz must be adapted for Hölder instead of Lipschitz, i.e. for $\gamma < 1.$

For 2., there is LipschitzOnWith.extend_real, which does not require the set A to be dense, but $\gamma = 1$ and $E = \mathbb{R}$. Also, there is DenseInducing.continuous_extend which gives a condition when a function can continuously be extended. (It needs a DenseInducing function, which in our case is $i : A \to D, x \mapsto x$.)

I:gauss Lemma 4.3. For $x \in \mathbb{R}$, let

$$\lfloor x \rfloor := \max\{n \in \mathbb{N} : n \le x\}.$$

The following holds:

- 1. $0 \le x |x| < 1;$
- 2. If $|x y| \le 1$, then $|\lfloor x \rfloor \lfloor y \rfloor| \le 1$.
- $3. |2\lfloor x \rfloor \lfloor 2x \rfloor| \le 1.$

Proof. 1. The first inequality is clear that $\lfloor x \rfloor$ is defined as a maximum over a set of numbers bounded above by x. The second inequality holds since otherwise we would have $\lfloor x \rfloor + 1 \leq x$, in contradiction to the definition of $\lfloor x \rfloor$.

2. Without loss of generality, assume that $y \leq x$ (which implies that $\lfloor y \rfloor \leq \lfloor x \rfloor$). The proof is by contradition, so assume that $\lfloor x \rfloor - \lfloor y \rfloor > 1$. So, we find $n := \lfloor x \rfloor \in \mathbb{N}$ such that $y < n-1 < n \leq x$. This means that x - y > n - (n-1) = 1, in contradiction to $|x - y| \leq 1$.

3. If $x - \lfloor x \rfloor < 1/2$, then $2x - 2\lfloor x \rfloor < 1$, which implies that $\lfloor 2x \rfloor = 2\lfloor x \rfloor$. Last, if $1/2 \le x - \lfloor x \rfloor < 1$, then $1 \le 2x - 2\lfloor x \rfloor < 2$, so $\lfloor 2x \rfloor = 2\lfloor x \rfloor + 1$ and the result follows.

Lemma 4.4. Let $I = [0, 1]^d$ and $|s - t| := \max_{i=1,...,d} |s_i - t_i|$ for $s, t \in I$. Let

- $D_n := \{0, 1, ..., 2^n\}^n \cdot 2^{-n} \subseteq I \text{ for } n = 0, 1, ..., \text{ and } D = \bigcup_{n=0}^{\infty} D_n;$
- $m \in \mathbb{N}$ and $s, t \in D$ with $|t s| \leq 2^{-m}$.

Then, there is $n \ge m$ and $s_m, ..., s_n, t_m, ..., t_n$ such that

- 1. $s_k, t_k \in D_k$ with $|s s_k|, |t t_k| \le 2^{-k}$ for all k = m, ..., n
- 2. $|s_k s_{k-1}|, |t_k t_{k-1}| \le 2^{-k}$,
- 3. $|t_m s_m| \le 2^{-m}$,
- 4. $s_n = s, t_n = t$.

Proof. Since $s, t \in D = \bigcup_n D_n$, and $D_n \subseteq D_m$ for $n \ge m$, there is some $n \ge m$ with $s, t \in D_n$. For $k \in m, ..., n$, we set

$$s_k := \lfloor s2^k \rfloor 2^{-k}, \qquad t_k := \lfloor t2^k \rfloor 2^{-k} \in D_k.$$

1. Since $|x - \lfloor x \rfloor| \le 1$ for all $x \in \mathbb{R}^d$ by Lemma ??.1, we have that

$$|s - s_k| = 2^{-k} |s2^k - \lfloor s2^k \rfloor| \le 2^{-k}, \qquad k = m, ..., n.$$

2. Using Lemma ??.3, write

$$|s_k - s_{k-1}| = 2^{-k} |\lfloor 2s2^{k-1} \rfloor - 2\lfloor s2^{k-1} \rfloor| \le 2^{-k}.$$

3. Since $|t-s| \leq 2^{-m}$, we have $|2^m t - s^m s| \leq 1$, so by Lemma ??.2

$$|t_m - s_m| = 2^{-m} |\lfloor t 2^m \rfloor - \lfloor s 2^m \rfloor| \le 2^{-m}.$$

4. We have
$$s2^n, t2^n \in \mathbb{Z}^d$$
 since $s, t \in D_n$, so $s_n = 2^{-n} \lfloor s2^n \rfloor = 2^{-n} s2^n = s$ and $t_n = t$.

rem1 Remark 4.5. Assume that $r(x_s, x_t) \leq 2^{-\gamma k}$ for all s, t with $|t - s| = 2^{-k}$ for $k \geq m$. Then, for some $s, t \in D$ with $|t - s| \leq 2^{-m}$, with s_k, t_k as in the above result and the triangle inequality,

$$t = t_n = s_n + \left(\sum_{k=m+1}^n t_k - t_{k-1} - (s_k - s_{k-1})\right) + t_m - s_m,$$

$$r(x_t, x_s) \le \left(\sum_{k=m+1}^n r(x_{t_k}, x_{t_{k-1}}) + r(x_{s_k}, x_{s_{k-1}})\right) + r(x_{t_m}, x_{s_m})$$

$$\le 2\sum_{k=m}^n 2^{-\gamma k} \le \frac{1}{1-2^{-\gamma}} 2^{-\gamma m}.$$

The proof of the continuity theorem follows the version in [?].

T:kolchen Theorem 2 (Continuous version; Kolmogorov, Chentsov). For some $d \in \mathbb{N}$ and $\sigma_1, \tau_1, ..., \sigma_d, \tau_d > 0$, let $I = \prod_{i=1}^d [\sigma_i, \tau_i]$, and $X = (X_t)_{t \in I}$ an E-valued stochastic process. Assume that there are $\alpha, \beta, C > 0$ with

$$\mathbf{E}[r(X_s, X_t)^{\alpha}] \le C|t-s|^{d+\beta}, \qquad 0 \le s, t \le \tau$$

There there exists a version $Y = (Y_t)_{t \in I}$ of X such that, for some random variables H > 0 and $K < \infty$,

$$\mathbf{P}\Big(\sup_{s\neq t, |t-s|\leq H} r(Y_s, Y_t)/|t-s|^{\gamma} \leq K\Big) = 1,$$

for every $\gamma \in (0, \beta/\alpha)$. In particular, Y almost surely is locally Hölder of all orders $\gamma \in (0, \beta/\alpha)$, and has continuous paths.

Proof. It suffices to show the assertion for $I = [0, 1]^d$. The general case then follows by some scaling of I. We consider the set of times

$$D_n := \{0, 1, ..., 2^n\}^n \cdot 2^{-n}$$

for $n = 0, 1, ..., D = \bigcup_{n=0}^{\infty} D_n$. Using the Markov inequality, we write for any $n \in \mathbb{N}$ (note that $|\{s, t \in D_n, |t-s| = 2^{-n}\}| \leq d2^{nd}$), using Lemma ??,

$$\mathbf{P}\left(\sup_{s,t\in D_n,|t-s|=2^{-n}}r(X_s,X_t)\geq 2^{-\gamma n}\right)\leq 2^{\gamma\alpha n}\mathbf{E}\left[\sup_{s,t\in D_n,|t-s|=2^{-n}}r(X_s,X_t)^{\alpha}\right]\\ \leq \sum_{s,t\in D_n,|t-s|=2^{-n}}2^{\gamma\alpha n}\mathbf{E}[r(X_t,X_s)^{\alpha}]\leq Cd2^{nd}2^{\gamma\alpha n}2^{-(d+\beta)n}=Cd2^{(\gamma\alpha-\beta)n},$$

and we see that the right hand side is summable. By the Borel-Cantelli Lemma,

$$N := \max\left\{n : \sup_{s,t \in D_n, |t-s|=2^{-n}} r(X_s, X_t) \ge 2^{-\gamma n}\right\} + 1$$

is finite, almost surely. From this and Remark ??, we conclude with $C' = \frac{1}{1-2^{-\gamma}}$,

$$\sup_{s,t\in D, s\neq t, |t-s|\leq 2^{-N}} r(X_s, X_t) \leq \sup_{m\geq N} \left(\sup_{s,t\in D, |t-s|\leq 2^{-m}} r(X_s, X_t) \right) \leq C' \sup_{m\geq N} 2^{-\gamma m} = C' 2^{-\gamma N}.$$

In other words, we see with $H = 2^{-N}$ and $K = C2^{-\gamma N}$,

$$\mathbf{P}\Big[\sup_{s,t\in D, s\neq t, |t-s|\leq H} r(X_s, X_t)/|t-s|^{\gamma} \leq K\Big] = 1,$$

i.e. X is locally Hölder- γ on D.

With this and Lemma ??.2, we can extend X Hölder-continuously on I, and call this extension $Y = (Y_t)_{t \in I}$. In order to show that Y is a modification of X, fix $t \in I$ and consider a sequence $t_1, t_2, \ldots \in D$ with $t_n \to t$ as $n \to \infty$. Then, for all $\varepsilon > 0$,

$$\mathbf{P}(r(X_{t_n}, X_t) > \varepsilon) \le \mathbf{E}[r(X_{t_n}, X_t)^{\alpha}] / \varepsilon^{\alpha} \xrightarrow{n \to \infty} 0,$$

i.e. $X_{t_n} \xrightarrow{n \to \infty} X_t$. Moreover, due to continuity of Y, we have $Y_{t_n} \xrightarrow{n \to \infty} f_s Y_t$. In particular, since $X_{t_n} = Y_{t_n}$ for all n, we have $\mathbf{P}(X_t = Y_t) = 1$ by Lemma ??, which concludes the proof. \Box

Lemma 4.6. Let (I, q) and (E, r) be metric spaces, and $f : I \to E$. Moreover, let $J \subseteq I$ be finite, $a, b, c \in \mathbb{R}_+$ with $a \ge 1$ and $n \in \{1, 2, ...\}$ such that $|J| \le ba^n$. Then, there is $K \subseteq J^2$ such that

$$|K| \le a|J|, \tag{2} eq:chain1$$

$$(s,t) \in K \Rightarrow q(s,t) \le cn,$$
 (3) eq:chain2

$$\sup_{s,t \in J, q(s,t) \le c} |f(t) - f(s)| \le 2 \sup_{(s,t) \in K} |f(s) - f(t)|.$$
(4) eq:chain3

Proof. Start with $V_1 = J$ and some $t_1 \in V_1$. We iteratively construct tuples $(V_\ell, t_\ell \in I, r_\ell \in \{1, ..., d\}, B_\ell \subseteq V_\ell, C_\ell \subseteq V_\ell, K_\ell \subseteq V_\ell^2)$ such that $V_{\ell+1} = V_\ell \setminus B_\ell$ (hence, $\ell \mapsto V_\ell$ is decreasing), $t_\ell \in V_\ell$ is some arbitrary element, and $(r_\ell, B_\ell, C_\ell, K_\ell)$ are given by first finding $r_\ell \in \{1, 2, ...\}$ minimal with

$$|C_{\ell}| \leq ba^{r_{\ell}}$$
 for $C_{\ell} := \{s \in V_{\ell} : q(s, t_{\ell}) \leq r_{\ell}c\}$

(since $|V_{\ell}| \leq ba^d$, this r_{ℓ} exists uniquely) and

$$B_{\ell} := \{ s \in V_{\ell} : r(s, t_{\ell}) \le (r_{\ell} - 1)c \} \subseteq C_{\ell}, \qquad K_{\ell} := \{ t_{\ell} \} \times C_{\ell}.$$

Note that this implies

$$|B_{\ell}| \ge ba^{r_{\ell}-1}, \qquad |K_{\ell}| = |C_{\ell}| \le ba^{r_{\ell}}$$

by definition of r_{ℓ} , and since $t_{\ell} \in B_{\ell}$ in all cases. We continue this construction until $V_m = \emptyset$. We claim that

$$K := \bigcup_{\ell=1}^{m} K_{\ell} = \{ (t_{\ell}, s) : s \in V_{\ell}, q(t_{\ell}, s) \le cr_{\ell} \text{ for some } \ell = 1, 2, \dots \}$$

satisfies (??), (??) and (??). In order to show (??), we have, since B_1, B_2, \ldots are disjoint,

$$\sum_{\ell} ba^{r_{\ell}-1} \leq \sum_{\ell} |B_{\ell}| \leq |J|.$$

Hence, (??) follows from

$$|K| \le \sum_{\ell} |K_{\ell}| = \sum_{\ell} |C_{\ell}| \le \sum_{\ell} ba^{r_{\ell}} \le a|J|.$$

For (??), we have for $(t_{\ell}, s) \in K_{\ell} \subseteq K$,

$$q(t_{\ell}, s) \le cr_{\ell} \le cd.$$

Last, for (??), consider $(s,t) \in J$ with $q(s,t) \leq c$. (Recall that $\ell \mapsto V_{\ell}$ is decreasing with $V_1 = J$ and $V_m = \emptyset$.) Find ℓ maximal with $s, t \in V_{\ell}$. Assume wlog that $s \notin V_{\ell+1}$, which implies $s \in B_{\ell}$ (since $V_{\ell+1} = V_{\ell} \setminus B_{\ell}$), which further implies $q(s,t_{\ell}) \leq (r_{\ell}-1)c$. (This implies $(t_{\ell},s) \in K_{\ell}$.) Since $q(s,t) \leq c$, this gives $q(t,t_{\ell}) \leq q(t,s) + q(s,t_{\ell}) \leq r_{\ell}c$, so $s, t \in C_{\ell}$. From here, $(t_{\ell},s), (t_{\ell},t) \in K_{\ell} \subseteq K$, hence

$$q(f(s), f(t)) \le q(f(s), f(t_{\ell})) + q(f(t_{\ell}), f(t)) \le 2 \sup_{s' \ t' \in K} q(f(s'), f(t'))$$

and we are done by taking $\sup_{s,t\in J,q(s,t)< c}$ on the left hand side.

Definition 4.7 (ε -cover). Let $\varepsilon > 0$ and (D, r) be some pseudometric space. A set $D' \subseteq D$ is said to be an ε -cover of D if $D = \bigcup_{x \in D'} B_{\varepsilon}(x)$. It is called minimal, if $D' \setminus \{x\}$ is no ε -cover for all $x \in D'$.

Lemma 4.8. Let $\varepsilon > 0$, (D, r) be some pseudometric space and $D_{\varepsilon} \subseteq D$ a minimal ε -cover of D. In addition, for $n = 1, 2, ..., set D_n := D_{2^{-n}}$.

- 1. For any $x \in D$ there is $x' \in D_{\varepsilon}$ such that $r(x, x') < \varepsilon$.
- 2. If $x \in D_{\varepsilon}$ is not isolated in D, there is $\varepsilon > 0$ and $x' \neq x$ with $x' \in D_{\varepsilon}$ and $r(x, x') < 3\varepsilon$.
- 3. Let $m \leq k$ and $x \in D_k$. Then, there is a sequence $x_k := x \in D_k, x_{k-1} \in D_{k-1}, ..., x_{m+1} \in D_{m+1}$ with $r(x_\ell, x_{\ell+1}) < 2^{-\ell-1}$ for $\ell = k, ..., m$.
- 4. Let $m \leq k, \ell, x \in D_k, y \in D_\ell$ with $r(x, y) < 2^{-m}$. Then, there are sequences $x_k := x, x_{k-1} \in D_{k-1}, ..., x_m \in D_m$ and $y_\ell := y, y_{\ell-1} \in D_{\ell-1}, ..., y_m \in D_m$ as in 2. with $r(x_m, y_m) < 3 \cdot 2^{-m}$.

Proof. 1. This follows from the definition of a ε -cover.

2. Since x is not isolated, there is $y \in D$ with $r(x, y) \in (\varepsilon, 2\varepsilon)$ for some $\varepsilon > 0$. Let $x' \in D_{\varepsilon}$ with $r(x', y) < \varepsilon$, which exists by 1. Then, $r(x, x') \le r(x, y) + r(y, x') < 3\varepsilon$.

3. As 1. shows, there is $x_{k+1} \in D_{k+1}$ with $r(x, x_{k+1}) < 2^{-k-1}$. The assertion follows inductively. 4. Using the triangle inequality,

$$r(x_m, y_m) \le r(x, y) + \sum_{i=m}^{k-1} r(x_i, x_{i+1}) + \sum_{i=m}^{\ell-1} r(y_i, y_{i+1})$$

$$< 2^{-m} + \sum_{i=m}^{\infty} 2^{-i-1} + \sum_{i=m}^{\infty} 2^{-i-1} = 3 \cdot 2^{-m}$$

Remark 4.9. Ok, this is simple, but, for d > 0

$$\sum_{j=0}^{n} 2^{dj} = \frac{2^{d(n+1)} - 1}{2^d - 1} \le \frac{2^d}{2^d - 1} 2^{dn}.$$

T:kolchen_general

Theorem 3 (Continuous version; Kolmogorov, Chentsov). Let (I,q) be a compact metric space and for $\varepsilon > 0$, let I_{ε} be a finite ε -cover of I. Assume that, for some $d \in \{1, 2, ...\}$, we have $c_1 > 0$ with

$$|I_{\varepsilon}| \le c_1 \varepsilon^{-d}$$

for ε small enough. Assume that $X = (X_t)_{t \in I}$ is an E-valued stochastic process and there are $\alpha, \beta, c_2 > 0$ with

$$\mathbf{E}[r(X_s, X_t)^{\alpha}] \le c_2 q(s, t)^{d+\beta}, \qquad s, t \in I.$$

Then, there exists a version $Y = (Y_t)_{t \in I}$ of X such that, for some random variables H > 0 and $K < \infty$,

$$\mathbf{P}\Big(\sup_{s\neq t,q(s,t)\leq H} r(Y_s,Y_t)/q(s,t)^{\gamma}\leq K\Big)=1,$$

for every $\gamma \in (0, \beta/\alpha)$. In particular, Y almost surely is locally Hölder of all orders $\gamma \in (0, \beta/\alpha)$, and has continuous paths.

Proof. With a slight abuse of notation, we set $I_n := I_{2^{-n}}$. (So, $|I_n| \leq c_1 2^{dn}$.) In addition, $J_n := \bigcup_{j=0}^n I_j$, hence $|J_n| \leq c_1 \sum_{i=0}^n 2^{di} \leq c_3 2^{dn}$ with $c_3 = \frac{2^d}{2^d-1}c_1$. We use Lemma ?? with $J = J_n$, $a = 2^d$, $b = c_3$, $c = 2^{-n}$ and n = n. This gives some $K_n \subseteq J_n^2$ with $|K_n| \leq c_3 2^d 2^{dn}$ such that $(s,t) \in K_n \Rightarrow q(s,t) \leq n \cdot 2^{-n}$ and

$$\sup_{s,t\in J_n,q(s,t)\leq 2^{-n}} r(X_s, X_t) \leq 2 \sup_{(s,t)\in K_n} r(X_s, X_t).$$

Using the Markov inequality, we write for any $n \in \mathbb{N}$, using Lemma ??,

$$\begin{aligned} \mathbf{P}\Big(\sup_{s,t\in J_n,q(s,t)<2^{-n}}r(X_s,X_t)\geq 2^{-\gamma n}\Big) &\leq \mathbf{P}\Big(2\sup_{(s,t)\in K_n}r(X_s,X_t)\geq 2^{-\gamma n}\Big) \\ &= \mathbf{P}\Big(\sup_{(s,t)\in K_n}r(X_s,X_t)^{\alpha}\geq 2^{-\alpha}2^{-\gamma\alpha n}\Big)\leq 2^{\alpha}2^{\gamma\alpha n}\mathbf{E}\Big[\sup_{(s,t)\in K_n}r(X_s,X_t)^{\alpha}\Big] \\ &\leq \sum_{(s,t)\in K_n}2^{\alpha}2^{\gamma\alpha n}\mathbf{E}[r(X_s,X_t)^{\alpha}]\leq c_32^d2^{nd}2^{\alpha}2^{\gamma\alpha n}c_2(3n\cdot2^{-n})^{d+\beta}=cn^{d+\beta}2^{(\gamma\alpha-\beta)n} \end{aligned}$$

with $c = c_3 2^d c_2$. So, we see that the right hand side is summable. By the Borel-Cantelli Lemma,

$$N := \max\left\{n : \sup_{s,t \in J_n, q(s,t) < 2^{-n}} r(X_s, X_t) \ge 2^{-\gamma n}\right\} + 1$$

is finite, almost surely. We set $J := \bigcup_n J_n$ and $H_m := 2^{-(N+m)}$ (with $H_m > 0$, almost surely). For $s \in J_k$ and k > N + m, let $s_k := s, s_{k-1}, ..., s_{N+m}$ be as in Lemma xxx, and analogously for $t \in K_\ell$ with $\ell > N + m$. From this and Remark ??, we conclude with $c' = 1 + \frac{2}{1-2-\gamma}$ and

$$\begin{split} \sup_{s,t\in J,q(s,t)\leq H_m} r(X_s, X_t) \\ &\leq \sup_{k,\ell\geq N+m} \sup_{s\in K_k, t\in K_\ell, q(s,t)\leq H_m} r(x_{s_{N+m}}, y_{t_{N+m}}) + \sum_{i=N+m}^{k-1} r(X_{s_i}, X_{s_{i+1}}) + \sum_{i=N+m}^{\ell-1} r(X_{t_i}, X_{t_{i+1}}) \\ &\leq \sup_{s,t\in J_{N+m}, q(s,t)\leq 3\cdot H_m} r(X_s, X_t) + 2^{-\gamma N} + \sum_{i=N}^{\infty} 2^{-\gamma i} + \sum_{i=N}^{\infty} 2^{-\gamma i} = c' 2^{-\gamma N}. \end{split}$$

In other words, we see with $H = 3 \cdot 2^{-N}$ and $K = c' 2^{-\gamma N}$,

$$\mathbf{P}\Big[\sup_{s,t\in J, s\neq t, q(s,t)\leq H} r(X_s, X_t)/q(s,t)^{\gamma} \leq K\Big] = 1,$$

i.e. X is locally Hölder- γ on K.

Then,

$$\sup \left\{ \frac{r(X_{s}, X_{t})}{q(s, t)^{\gamma}} : s, t \in J, q(s, t) \leq 3 \cdot 2^{-N} \right\}$$

$$= \sup_{m=0,1,2,\dots} \sup \left\{ \frac{r(X_{s}, X_{t})}{q(s, t)^{\gamma}} : s, t \in J, H_{m+1} < q(s, t) \leq H_{m} \right\}$$

$$\leq \sup_{m=0,1,\dots} 2^{(N+m+1)\gamma} \sup \left\{ r(X_{s}, X_{t}) : s, t \in J, q(s, t) \leq H_{m} \right\}$$

$$\leq \sup_{m=0,1,\dots} 2^{(N+m+1)\gamma} c' 2^{-\gamma(N+m)}$$

$$= 2^{\gamma} c'.$$

With this and Lemma ??.2, we can extend X Hölder-continuously on I, and call this extension $Y = (Y_t)_{t \in I}$. In order to show that Y is a modification of X, fix $t \in I$ and consider a sequence $t_1, t_2, \ldots \in D$ with $t_n \to t$ as $n \to \infty$. Then, for all $\varepsilon > 0$,

$$\mathbf{P}(r(X_{t_n}, X_t) > \varepsilon) \le \mathbf{E}[r(X_{t_n}, X_t)^{\alpha}] / \varepsilon^{\alpha} \xrightarrow{n \to \infty} 0,$$

i.e. $X_{t_n} \xrightarrow{n \to \infty}_p X_t$. Moreover, due to continuity of Y, we have $Y_{t_n} \xrightarrow{n \to \infty}_{fs} Y_t$. In particular, since $X_{t_n} = Y_{t_n}$ for all n, we have $\mathbf{P}(X_t = Y_t) = 1$ by Lemma ??, which concludes the proof. \Box

References

- [EK86] S.N. Ethier and T.G. Kurtz. Markov Processes. Characterization and Convergence. John Wiley, New York, 1986.
- [KS91] Ioannis Karatzas and Steven Shreve. Brownian motion and stochastic calculus, volume 113. Springer Science & Business Media, 1991.