

# Formalizing perfectoid spaces

## Abstract

Perfectoid spaces are sophisticated objects in arithmetic geometry introduced by Peter Scholze in 2012. We formalized enough definitions and theorems in topology, algebra and geometry to define perfectoid spaces in the Lean theorem prover. This experiment confirms that a proof assistant can handle complexity in that direction, which is rather different from formalizing a long proof about simple objects. It also confirms that mathematicians with no computer science training can become proficient users of a proof assistant in a relatively short period of time. Finally, we observe that formalising a piece of mathematics that is a trending topic boosts the visibility of proof assistants amongst pure mathematicians.

**Keywords** formal proof, algebraic geometry, Lean

## 1 Introduction

In 2012, Peter Scholze defined the notion of a perfectoid space, and used it to prove new cases of the weight-monodromy conjecture, an extremely important conjecture in modern arithmetic geometry. This original application of the theory was based on a key theorem of Scholze called the *tilting correspondence*, relating perfectoid spaces in characteristic zero to those in positive characteristic. Over the next few years, many other applications appeared, culminating in some spectacular applications to the Langlands programme. Scholze was awarded the Fields Medal in 2018 for his work. See [21, 26] for far more thorough explanations of how Scholze’s ideas have changed modern mathematics.

With current technology, it would take many person-decades to formalise Scholze’s results. Indeed, even *stating* Scholze’s theorems would be an achievement. Before that, one has of course to formalise the definition of a perfectoid space, and this is what we have done, using the Lean theorem prover. This requires formalizing many intermediate definitions and proving many theorems, and was done without adding any axioms to Lean’s logic. Formalizing the definition of such sophisticated objects is rather different from formalizing a complex proof about very simple objects, as in [11]. Contemporary mathematics is also formalised in [7, 12], but those papers focus on elementary objects.

The goal of this text is to report on the formalization process, highlighting our efforts to bridge the gap between the formal and informal stories. The level of details decreases quickly along the paper, mostly because the involved mathematics becomes too technical for the format constraints,

but also because the relatively elementary parts of the story probably are interesting to more people.

As far as we know, no other formalisation of perfectoid spaces has been attempted in any theorem prover. A priori it could be done, although it seems more difficult without dependent types. For instance, sheaves are functions associating a ring to each open subset of a topological space, an idea which is naturally expressed as a dependent type.

Our main source for the classical part of the story is Bourbaki’s *Elements of mathematics*. More recent sources all seem to focus on special cases that are not general enough for our purposes, and defer to Bourbaki for the general case. Of course Bourbaki’s careful style is also very valuable. We hope our readers won’t be misled by the fact that we do point out a couple of arguments that we could improve on. That’s simply because repeating everywhere else that Bourbaki was spot on does not make a very interesting story.

Our main source for the recent part of the story is Wedhorn’s lecture notes [25], with occasional need to go back to the original sources by Huber and Scholze [14, 15, 23, 24].

## 2 Background and overview

A perfectoid space is an adic space satisfying some properties. The history of adic spaces goes back to the basic question of how to do analysis over complete fields such as the field  $\mathbb{Q}_p$  of  $p$ -adic numbers, and other fields of interest to number theorists. The classical theory of analysis of one complex variable has a local definition of differentiation, with powerful consequences such as the theorem that a differentiable function on the complex plane has a power series expansion which converges everywhere. Whilst there was clearly some  $p$ -adic analogue of the theory (for example  $p$ -adic exponential and logarithm functions shared many properties with their classical counterparts), the foundations of the subject remained elusive for the simple reason that  $\mathbb{Q}_p$  is totally disconnected, and there are plenty of examples of differentiable functions which were locally constant but not constant. To give a specific example, consider the function on the closed unit disc  $\mathbb{Z}_p$  defined as the characteristic function of the open unit disc (so 1 on the open disc and 0 elsewhere). It is locally constant (as both the open unit disc and its complement are open, due to the non-Archimedean nature of the metric), and hence satisfies the naive definition of differentiability, however it has no global power series expansion.

In the late 1960s, Tate proposed a foundation to the theory which solved these problems, namely the theory of rigid analytic spaces. Tate starts with the observation that each point  $a$  in  $\mathbb{Z}_p$  yields a maximal ideal of the ring  $\mathbb{Q}_p\langle X \rangle$  of power series which converge on  $\mathbb{Z}_p$ , namely the kernel of

the ring homomorphism  $\mathbb{Q}_p\langle X \rangle \rightarrow \mathbb{Q}_p$  sending  $f$  to  $f(a)$ . Tate's replacement for the closed unit disc is the collection of all maximal ideals of this ring. Using "spaces" formed by maximal ideals of appropriate rings, Tate developed a  $p$ -adic theory analogous to the theory of real or complex manifolds. This theory had all the right properties, but the underlying "spaces" were no longer topological spaces, they were spaces equipped with a Grothendieck topology. Tate's resolution of the problem with the characteristic function of the open unit disc is that the cover of the rigid analytic closed unit disc by the open disc and its complement is not admissible for his Grothendieck topology, even though both sets are open.

However, the use of a Grothendieck topology was inconvenient. In the 1990s, Berkovich and Huber proposed variants of the theory. The maximal ideals in Tate's algebras  $A$  gave rise to maps from  $A$  to finite extensions of  $\mathbb{Q}_p$ , and hence to valuations on  $A$ . So we can add even more points to our spaces by considering more valuations on these rings. Berkovich considered real-valued valuations, but Huber allowed his valuations to take values in certain totally ordered monoids. Then many extra points appear on or near the boundary of the open unit disc, with the result that the function defined to be 1 on the open disc and 0 outside it was no longer a continuous function; the complement of the open disc was no longer open. For a vivid description of the points that get added to the closed unit disc over a complete algebraically closed field, see [23, Example 2.20].

Huber called his new objects *adic spaces*, and the majority of our work is a formalisation of their definition. Adic spaces represent a decisive generalisation of Tate's theory of rigid analytic spaces, because finiteness constraints (such as finite-dimensionality of the objects in question) are embedded in Tate's foundations, but are not necessary in the theory of Berkovich or adic spaces. This was crucial for Scholze, whose perfectoid spaces are built from rings which almost never satisfy these constraints.

Adic spaces have more structure than a topology. They are endowed with a sheaf of complete topological rings, and equivalence classes of valuations on the stalks of this sheaf. In particular, this structure associates to each open subset  $U$  in our space a ring equipped with a topology that is compatible with its ring structure, and is complete in a sense that generalizes the definition of complete metric spaces. This ring is meant to describe analytic functions on  $U$ .

In order to motivate the introduction of completeness in this story, let us evoke how to go from a polynomial ring  $k[X]$  to the corresponding power series ring  $k[[X]]$  from a topological point of view. One can define a topology on  $k[X]$ , which is compatible with its ring structure, and such that neighborhoods of 0 are subsets containing the ideal  $X^n k[X]$  for some natural number  $n$ . Then the partial sums  $S_N$  of a power series can be seen as a sequence in  $k[X]$  which is Cauchy for this topology: the difference  $S_N - S_M$  belongs to an arbitrarily small neighborhood of zero as  $N$  and  $M$  tend

to infinity. One can then view  $k[[X]]$  as a completion of  $k[X]$  where this sequence converges.

Adic spaces are constructed by gluing model spaces associated to certain kinds of topological rings. This is analogous to how manifolds are locally modelled on open subsets of affine spaces, and schemes are locally modelled on spectra of rings. But here the local models are more complicated to build, and *proving* all the axioms expected from the extra structure means proving a lot of theorems. This is the bulk of our formalization effort. It involves a subtle blend of general topology and commutative algebra.

All the story so far brings us to the 90s. Briefly, the motivation behind Scholze's perfectoid spaces is the following. Some of the simplest examples of maps between geometric objects are the unramified maps: maps which are locally isomorphisms. This geometric notion has an arithmetic counterpart; the extensions of local fields which are easiest to study are the unramified extensions. Again it was Tate who showed us another approach; if instead one considered so-called "deeply ramified" extensions, then they were so badly ramified everywhere that any finite extension of a deeply ramified extension was "almost unramified". This idea was taken further first by Faltings, and then by Scholze, who extended the theory to adic spaces: Scholze's perfectoid spaces can be thought of as the "deeply ramified" adic spaces. Their geometry is governed by their characteristic  $p$  "tilts", giving a profound link between geometry in characteristic  $p$  and characteristic 0. We won't give more details about this motivation here, since writing the definition of perfectoid spaces is actually pretty short once adic spaces are defined.

### 3 Lean and its mathematical library

Our formalization of perfectoid spaces uses the Lean proof assistant [8], developed principally by Leonardo de Moura since 2013 at Microsoft Research. Lean implements a version of the calculus of inductive constructions [6], with quotient types, non-cumulative universes, and proof irrelevance.

In 2017, almost all the mathematical content of Lean's core library was split off as a community curated library called `mathlib` [1], which is the basis for all mathematical developments in Lean, including our perfectoid spaces formalization. Initially developed mostly by Mario Carneiro and Johannes Hölzl, this library then gained many contributors, including many people coming from mathematics departments. Ultimately, everything described in this paper until Section 6 included should be merged into `mathlib`, leaving only very specialized material in the perfectoid spaces project. Some of our work was directly done inside `mathlib`, some was merged later, some is still to be merged.

The `mathlib` library takes inspiration from many great efforts in other proof assistants, notably Isabelle, Coq, and Metamath, but also has its own distinct flavor. Of course the most relevant comparison is with Mathematical Components

[19], since Lean's logical foundations are extremely close to those of Coq. On the mathematical side, the main difference is that `mathlib` does not insist on constructivity. This makes it very appealing to regular mathematicians. On the technical side, a very visible difference is `mathlib`'s heavy use of type classes instead of canonical structures as in [9]. Both these mechanisms allows to handle extra structures on types. For instance, for every type  $G$ , `add_group G` is a type packaging a composition law  $+$ , a term  $\theta$  of type  $G$  and properties like  $\forall g, g+\theta = g$  etc. The type family `add_group` is declared to be a *type class*. One can then declare *instances* of that class. They can be concrete instances, like a group structure on  $\mathbb{Z}$ , or composite instances, such as the instance creating a group structure on  $G \times H$  from a group structure on  $G$  and a group structure on  $H$ . After setting up such a class and some instances, one can declare that certain arguments of later definitions and lemmas should be automatically inferred by searching the instance database, this is the instance resolution procedure.

During the instance resolution procedure, Lean can only identify terms and types that are definitionally equal, it does not use theorems asserting equalities. This leads to encoding tricks that are already present in Coq<sup>1</sup> and Isabelle for instance, but are very counter-intuitive for mathematicians. For instance, a metric space structure on some  $X$  is usually defined as a package containing a distance function from  $X \times X$  to  $\mathbb{R}_{\geq 0}$  and a bunch of conditions on this function. Then one can define the topology associated to a metric space structure, and get access to theorems proved about general topological spaces. In `mathlib`, and some other libraries, a metric structure on  $X$  is<sup>2</sup> a package containing not only the distance function and its properties, but also a topology and a property asserting that this topology is compatible with the distance function. Of course there is a function which takes as input the traditional package and outputs this "augmented" package, so defining concrete instances is no harder. The difference matters with composite instances. Say we start with two metric spaces  $X$  and  $Y$ , and we want a metric structure on  $X \times Y$ . The traditional way would define a distance on  $X \times Y$ , check the axioms, and then, after this construction, prove a compatibility lemma saying the topology built from the product metric equals the product topology. But that equality will not be definitional, and, in particular, this could block further instance resolution. In the augmented packaging approach, building the product instance involves defining the product distance, feeding in the product topology, and using the mathematical content of the compatibility lemma to provide the compatibility part of the package. So the total amount of mathematical work is the same, but the metric topology on  $X \times Y$  is now definitionally equal to the product of the metric topologies of  $X$  and  $Y$ .

<sup>1</sup>see e.g. [9, Section 3.2]

<sup>2</sup>Actually what we describe is a simplified story.

## 4 Topology, filters and extensions

### 4.1 Topology without subsets

Topology in `mathlib` began when Johannes Hölzl ported his own Isabelle work, documented in [13]. The main traditional source for this is Bourbaki [3]. But the formal version goes quite a bit further in the systematic and functorial use of filters and order relations, as we will illustrate below.

Filters were introduced by Henri Cartan in order to unify various notions of limits in topology. They became standard in formalized libraries since [13], see [2, 22]. A filter on  $X$  is a set  $\mathcal{F}$  of subsets of  $X$  which contains  $X$ , is stable under pairwise intersection, and such that every subset containing an element of  $\mathcal{F}$  belongs to  $\mathcal{F}$ . The original definition was restricted to what we now call *proper* filters: those not containing the empty subset; the new definition is however more functorial and hence more convenient.

Every point  $x$  in a topological space  $X$  gives rise to the filter  $\mathcal{N}_x$  of neighborhoods of  $x$ . Complements of finite subsets in  $\mathbb{N}$  also form a filter  $\mathcal{F}_{\mathbb{N}}$ . In  $\mathbb{R}$  one can also consider  $\mathcal{N}_{+\infty}$  made of all subsets containing some  $[A, +\infty)$ , and similarly  $\mathcal{N}_{-\infty}$ .

Given a filter  $\mathcal{F}$  on  $X$  and a map  $f: X \rightarrow Y$ , the pushforward filter  $f_*\mathcal{F}$  on  $Y$  consists of subsets of  $Y$  whose preimage under  $f$  belong to  $\mathcal{F}$ . The pullback  $f^*\mathcal{G}$  of a filter  $\mathcal{G}$  on  $Y$  consists of subsets of  $X$  containing the preimage of an element of  $\mathcal{G}$ . For instance, if  $X$  is equipped with a topology induced from  $Y$  by  $f$  then  $\mathcal{N}_x = f^*\mathcal{N}_{f(x)}$ . Those operations are functorial:  $(f \circ g)_* = f_* \circ g_*$  and  $(f \circ g)^* = g^* \circ f^*$ .

A filter  $\mathcal{F}$  is *finer* than another one  $\mathcal{F}'$  (on the same  $X$ ) if  $\mathcal{F}' \subseteq \mathcal{F}$ . We denote this order relation by  $\mathcal{F} \leq \mathcal{F}'$ . Both mapping operations  $f_*$  and  $f^*$  are order preserving, and they form a Galois connection:  $f_*\mathcal{F} \leq \mathcal{G} \iff \mathcal{F} \leq f^*\mathcal{G}$ .

A filter  $\mathcal{G}$  on some topological space  $Y$  *converges* to a point  $y \in Y$  if  $\mathcal{G} \leq \mathcal{N}_y$ . A function  $f$  from  $X$  to  $Y$ , *tends* to a point  $y$  with respect to a filter  $\mathcal{F}$  on  $X$  if  $f_*\mathcal{F}$  converges to  $y$ . As announced this allows to unify the definition of limit of a sequence  $u: \mathbb{N} \rightarrow Y$ , a limit of a function  $f: X \rightarrow Y$  at some point  $x$  in a topological space  $X$ , a limit of  $f: \mathbb{R} \rightarrow Y$  at  $\pm\infty$ , using filters  $\mathcal{F}_{\mathbb{N}}$ ,  $\mathcal{N}_x$  and  $\mathcal{N}_{\pm\infty}$  respectively. But, as noted in [13], one can go one step further, consider any filter  $\mathcal{F}'$  on  $Y$ , and say that  $f$  tends to  $\mathcal{F}'$  with respect to  $\mathcal{F}$  if  $f_*\mathcal{F} \leq \mathcal{F}'$ .

Unpacking the definitions, one can check for instance that  $f$  tends to  $y$  with respect to  $\mathcal{N}_x$  means that the preimage  $f^{-1}(V)$  of any neighborhood  $V$  of  $y$  is a neighborhood of  $x$ . But such an unpacking is often not needed, and avoiding it leads to very efficient proofs by computation (in the mathematical sense, not  $\lambda$ -calculus), especially when combined with a systematic use of lattice operations on the type of topologies or filters on a type. For instance, given topologies  $T$  and  $S$  on  $X$  and  $Y$ , the product topology on  $X \times Y$  is defined as  $\inf(\text{pr}_1^*T, \text{pr}_2^*S)$ , using the general notion of induced topology and infimum of topologies. Similarly, products of filters are defined by  $\mathcal{F} \times \mathcal{G} = \inf(\text{pr}_1^*\mathcal{F}, \text{pr}_2^*\mathcal{G})$ , and satisfy

$f_*(\mathcal{F} \times \mathcal{G}) \leq f_*\mathcal{F} \times f_*\mathcal{G}$ , and  $\mathcal{N}_{(x,y)} = \mathcal{N}_x \times \mathcal{N}_y$ . None of these formulas explicitly feature subsets, they operate at a higher level of abstraction. As an example of using them, let us look at our proof that if  $f : Z \rightarrow X$  and  $g : Z \rightarrow Y$  are continuous at  $z_0$  then so is  $h := (z \mapsto (f(z), g(z)))$ . In Lean code below,  $\text{map } h \mathcal{F}$  means  $h_*\mathcal{F}$ , and  $\times^f$  is product of filters.

```

lemma continuous_at.prod_mk {f : Z → X} {g : Z → Y}
{z₀ : Z} (h : continuous_at f z₀) (h' : continuous_at g z₀):
continuous_at (λ z, (f z, g z)) z₀ :=
calc map (λ z, (f z, g z)) (ℳ z₀)
  ≤ (map f ℳ z₀) ×f (map g ℳ z₀) : map_prod_mk _ _ _
  ... ≤ (ℳ f z₀) ×f (ℳ g z₀) : prod_mono h h'
  ... = ℳ (f z₀, g z₀) : nhds_prod_eq.symm

```

This style of proof was strongly advocated by Hölzl, who initiated that part of `mathlib`. It does not seem to be used in Isabelle or Coq. In addition to a couple of sporadic lemmas, like the above one, we had to develop the theory of filter bases, which is not so useful for the abstract story, but very convenient to define concrete instances, like the adic topology defined by an ideal  $I$  in a ring  $R$ . This  $I$ -adic topology is characterized by the fact that neighborhoods of zero are subsets containing a power of  $I$ .

## 4.2 Extension by continuity

Extension by continuity is an important operation of elementary topology, and it plays a crucial role in the completion of topological rings that we need in order to define the structure presheaf on adic spectra. The following is [3, Theorem I.8.5.1] (recall that a topological space is regular if every point has a basis of closed neighborhoods).

**Theorem 4.1.** *Let  $X$  be a topological space,  $A$  a dense subset of  $X$ , and  $f : A \rightarrow Y$  a mapping of  $A$  into a regular space  $Y$ . A necessary and sufficient condition for  $f$  to extend to a continuous mapping  $\tilde{f} : X \rightarrow Y$  is that, for each  $x \in X$ ,  $f(y)$  tends to a limit in  $Y$  when  $y$  tends to  $x$  while remaining in  $A$ . The continuous extension  $\tilde{f}$  of  $f$  to  $X$  is then unique.*

From a type theory point of view, we immediately spot a little difficulty here since we need every function to be total on its domain type. One way to fix this is to make the inclusion  $\text{map } \iota : A \rightarrow X$  more visible. It is actually already present since “ $y$  tends to  $x$  while remaining in  $A$ ” means that we use the filter  $\iota^*\mathcal{N}_x$ . So the extension problem becomes a factoring problem, i.e. finding  $\tilde{f}$  such that  $\tilde{f} \circ \iota = f$ .

This little perspective shift allows to consider the case where  $\iota$  is not injective. And we will use this case in the theory of uniform spaces (in particular topological groups and rings). We simply assume that  $A$  and  $X$  are topological spaces,  $\iota$  pulls back the topology of  $X$  onto the topology of  $A$ , and the image  $\iota(A)$  is dense in  $X$ .

Another issue related to totality of functions is that Theorem 4.1 is too global. We want a version that applies to

functions that are not everywhere continuous. The key example is inversion on a topological field. Inversion needs to be a total function so the inverse of zero has to be defined. Of course we don't expect inversion to be continuous at zero, but we still want to use some version of Theorem 4.1 in order to define inversion on  $\mathbb{R}$  out of inversion on  $\mathbb{Q}$  for instance. The traditional workaround is to apply the theorem to  $\mathbb{Q} \setminus \{0\}$  equipped with the induced topology. We could do that in Lean of course, but it creates some overhead which is more difficult to ignore than on paper, and tends to pile up.

One last formalization detail is that we do not want to carry around the limit assumption each time we mention the extension  $\tilde{f}$ . We want an extension operator  $E_\iota$  transforming functions from  $A$  to  $Y$  into functions from  $X$  to  $Y$ . Of course the result won't have any useful properties if  $f$  is awful. We can do that using the axiom of choice. The definition of  $E_\iota f(x)$  is: if there exists some  $y$  such that  $f_*\iota^*\mathcal{N}_x \leq \mathcal{N}_y$  then choose such a  $y$  as  $f(x)$ , otherwise choose any element of  $Y$ . Our final version of the extension theorem is then:

**Theorem 4.2.** *Let  $\iota : A \rightarrow X$  be a map between topological spaces whose image is dense, and which pulls back the topology of  $X$  to the topology of  $A$ . Let  $f$  be a map from  $A$  to a regular topological space  $Y$ .*

- If  $f$  is continuous at  $a$  then  $E_\iota f(\iota(a)) = f(a)$ .
- Let  $x$  be a point in  $X$ . If, for every  $x'$  in a neighborhood of  $x$ , there exists  $y$  in  $Y$  such that  $f_*\iota^*\mathcal{N}_{x'} \leq \mathcal{N}_y$  then  $E_\iota f$  is continuous at  $x$ .

Note that the proof is almost exactly the same as the proof of Theorem 4.1, which uses injectivity of  $\iota$  only for psychological purposes. And of course locality also comes essentially for free in the proof since continuity is a local property.

Theorems 4.1 and 4.2 have an obvious drawback: the existence of a limit assumption looks fairly difficult to check. In undergraduate accounts of this theorem, it is replaced by a uniform continuity assumption on  $f$  coupled with a completeness assumption on the target space  $Y$ . In such accounts both assumptions seem to require using metric spaces, which is too restrictive for our purposes. The next section addresses this problem by introducing uniform spaces. This will allow us to go beyond metric spaces, but won't solve the issue that uniform continuity is too much to ask for. For instance, even in the most elementary case of extending multiplication from  $\mathbb{Q} \times \mathbb{Q} \rightarrow \mathbb{Q}$  to  $\mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  we hit the problem that this map is not uniformly continuous. In this case it is locally uniformly continuous, but this won't happen for general topological rings. A very good introduction to the next two sections is [17], where Lewis explains his design decisions for the definition of  $\mathbb{Q}_p$ , with its valued field structure<sup>3</sup>. He is able to take several shortcuts that are not available to us in the more general setting of Huber rings and adic spaces.

<sup>3</sup>There is also a formalization of  $\mathbb{Q}_p$  in [20] from a purely algebraic point of view, which is irrelevant here

## 5 Completions

### 5.1 Uniform notions

Uniform structures are refinements of topological structures introduced by André Weil as a common generalization of metric spaces and topological groups, which allow to discuss uniform continuity (hence the name) and completeness. A *uniform structure* on  $X$  is a filter  $\mathcal{U}$  on  $X \times X$  satisfying some conditions. Elements of  $\mathcal{U}$  are called entourages. Two moving points  $x$  and  $y$  are declared arbitrarily close to each other if the pair  $(x, y)$  belongs to an arbitrary entourage. The first condition imposed on  $\mathcal{U}$  is that all entourages contain the diagonal. This is analogous to the reflexivity condition in the definition of a distance function. The other conditions are analogous to the symmetry and triangle inequality conditions. However, no analogue of the separation axiom  $d(x, y) = 0 \implies x = y$  is enforced, allowing for non separated spaces. One can prove that, given a uniform structure  $\mathcal{U}$ , there is a unique topology  $T_{\mathcal{U}}$  on  $X$  such that  $\mathcal{N}_x = (y \mapsto (x, y))^* \mathcal{U}$ . This topology is separated if and only if the diagonal is the only subset of  $X \times X$  contained in all entourages. In `mathlib`, the topology is actually part of the uniform structure, together with an axiom equivalent to the neighborhood formula. This is an encoding trick fully analogous to what we explained in Section 3.

Uniform structures can be pulled back: one can check that, for every map  $f : X \rightarrow Y$  and every uniform structure  $\mathcal{U}$  on  $Y$ , the filter  $(f \times f)^* \mathcal{U}$  is a uniform structure on  $X$  (however pushforward does not work in general, the analogue of the triangle inequality is not preserved). One can then define uniform continuity for maps  $f$  between uniform spaces  $X$  and  $Y$  by  $\mathcal{U}_X \leq (f \times f)^* \mathcal{U}_Y$ . Again this way of writing things is slightly more abstract than what is usually seen in textbooks. For instance, Bourbaki writes:  $f : X \rightarrow Y$  is uniformly continuous if, for each entourage  $V$  of  $Y$ , there is an entourage  $W$  of  $X$  such that the relation  $(x, x') \in W$  implies  $(f(x), f(x')) \in V$ . Bourbaki then writes that an immediate consequence of definitions is that uniformly continuous functions are continuous (for the underlying topologies). But thinking in more abstract terms actually allows to break the proofs into small lemmas that are needed anyway, and gives a computational proof which does not mention entourages at all. The lemmas assert that taking the topology underlying a uniform structure is order preserving and “commutes with pull-back” (the later being called the “commutativity property” until the end of this paragraph). The proof of continuity then becomes:  $T_{\mathcal{U}_X} \leq T_{(f \times f)^* \mathcal{U}_Y} = f^* T_{\mathcal{U}_Y}$  where the inequality comes from the order preserving property of  $T$  applied to the uniform continuity assumption, and the equality is the commutativity property. The resulting inequality  $T_{\mathcal{U}_X} \leq f^* T_{\mathcal{U}_Y}$  is equivalent to continuity of  $f$ . In `mathlib`, the commutativity property is definitional, by the encoding trick, so the proof does not even need to mention it, and ends

up being only twice as many characters long as the sentence “This follows immediately from definitions”.

To each metric space  $(X, d)$ , we can associate the uniform structure whose entourages are sets containing some  $\{(x, y) \mid d(x, y) \leq \varepsilon\}$  for some positive  $\varepsilon$ . Then the usual definition of uniform continuity is recovered. We include this example in order to relate uniform spaces to metric spaces, but it doesn't play any role in our formalization. A relevant example associates to each topological abelian group  $G$ , with subtraction map  $\delta : G \times G \rightarrow G$ , the uniform structure  $\delta^* \mathcal{N}_0$ . Here we assume  $G$  is commutative for simplicity. In our applications it will always be the underlying additive group of a topological ring.

Similarly, one can extend the definition of Cauchy sequences to topological groups. However this notion is not powerful enough when sequential continuity does not imply continuity. The solution here is to define a notion of Cauchy filters, not necessarily coming from sequences. A filter  $\mathcal{F}$  of  $X$  is Cauchy if it is proper and if  $\mathcal{F} \times \mathcal{F} \leq \mathcal{U}$ . This indeed generalizes the notion of Cauchy sequences: a sequence  $u : \mathbb{N} \rightarrow X$  into a metric space or a topological group is Cauchy if and only if the filter  $u_* \mathcal{F}_{\mathbb{N}}$  is Cauchy (in `mathlib`, Cauchy sequences are actually defined like this, but the elementary definition is then proved to be equivalent). By definition, a uniform space is complete if every Cauchy filter converges to some point. This is equivalent to the elementary notion in the case of metric spaces.

Uniform spaces, uniform continuous functions and Cauchy filters were formalized before in Isabelle/HOL and Coq [2, 22]. The Coq formalizations currently use a slightly awkward definition of uniform spaces, but [22] mentions plans to switch to the one explained above. As far as we know, everything from this point to the end of the paper has never been formalized before.

### 5.2 The completion functor

As explained in the introduction, completing topological rings (equipped with their uniform structure) is the required abstraction of moving from polynomials to general analytic functions. It happens that separated completions always exist for uniform spaces. More precisely, the inclusion functor from the category of complete separated uniform spaces into the category of all uniform spaces admits a left adjoint. The key part of that assertion is the following theorem:

**Theorem 5.1** ([3, Theorem II.3.7.3 and Definition II.3.7.4]). *For each uniform space  $X$ , there is a complete separated uniform space  $\hat{X}$  and a uniformly continuous map  $\iota_X : X \rightarrow \hat{X}$  with the following universal property: every uniformly continuous map  $f$  from  $X$  to a complete separated uniform space  $Y$  uniquely factors through  $\iota_X$ . The space  $\hat{X}$  is called the separated completion of  $X$ .*

One can apply this universal property to  $\iota_Y \circ f$  when  $Y$  is not complete and separated to lift  $f$  to a morphism

551  $\hat{f} : \hat{X} \rightarrow \hat{Y}$ . Uniqueness in the universal property then  
 552 proves this lifting operation respects composition, hence we  
 553 indeed have a completion functor.

554 Bourbaki constructs  $\hat{X}$  as the space of maximal<sup>4</sup> Cauchy  
 555 filters on  $X$ , and the map  $\iota_X$  sends  $x$  to  $\mathcal{N}_x$ . The first main  
 556 task is to endow this space with a uniform structure which  
 557  $\iota_X$  pulls back to  $\mathcal{U}_X$ , and prove that  $\iota_X$  has dense range. The  
 558 next task is to build factorizations. Of course one wants to  
 559 use extension by continuity, using completeness of the target  
 560 space to provide the necessary limits. But  $\iota_X$  cannot be  
 561 injective if  $X$  is not separated. Indeed any uniformly continuous  
 562 map into a separated space sends non-separated pairs of  
 563 point into the same point. So Bourbaki needs to first factor  
 564 through the  $\iota_X(X)$ , and then apply Theorem 4.1. With our  
 565 Theorem 4.2, this detour is not necessary.

566 In `mathlib`, we used a variation on this construction, sug-  
 567 gested in [16] and whose formalization was started by Hölzl.  
 568 One could wonder how to relate this construction to the  
 569 Bourbaki construction. More seriously, there are more ele-  
 570 mentary constructions of completions in the special case of  
 571 metric spaces. The `mathlib` library constructs real numbers  
 572 and  $p$ -adic numbers using equivalence classes of Cauchy  
 573 sequences of rational numbers. We certainly want to make  
 574 sure these are isomorphic to the our general completions, if  
 575 only as a sanity check (note Bourbaki solves this problem  
 576 by avoiding any mention of the elementary approach). We  
 577 introduce the following definition. Given a uniform space  $X$ ,  
 578 we say that  $(Y, \iota : X \rightarrow Y)$  is an *abstract completion of  $X$*  if  $Y$   
 579 is a complete separated space,  $\iota$  has dense image and pulls  
 580 back the uniform structure of  $Y$  onto the uniform structure  
 581 of  $X$ . We then prove that this property is sufficient to ensure  
 582 the expected universal property, without any reference to  
 583 the construction of  $Y$ , which is left completely unspecified.  
 584 The usual abstract arguments then gives an isomorphism  
 585 between any two abstract completions of a given uniform  
 586 space. We can then check that both our general construction  
 587 and the Cauchy sequences completion of rational numbers  
 588 are abstract completions, hence isomorphic.

### 589 5.3 Completing groups and rings

591 As explained above, topological groups have a natural uni-  
 592 form structure. Mathematically it sounds natural to define a  
 593 type class instance of `uniform_space`  $G$  from an instance of  
 594 `topological_add_group`  $G$ . One can then use the comple-  
 595 tion construction from the preceding section. Extending the  
 596 group operations to the completion  $\hat{G}$  is immediate using  
 597 the completion functor, since these operations are uniformly  
 598 continuous. The extensions are uniformly continuous hence  
 599 continuous, and  $\hat{G}$  is also a topological group. But this leads  
 600 to a subtle issue: we now have two different uniform struc-  
 601 tures on  $\hat{G}$ . The first one comes from the completion functor  
 602 composed with the forgetful functor from uniform spaces  
 603

604 <sup>4</sup>Bourbaki writes “minimal” because his order on filters is opposite to ours.

606 to topological spaces. The second one comes from the topo-  
 607 logical group structure. Those two uniform structures are  
 608 equal, but this is a theorem which is not obvious in any  
 609 way. Bourbaki does prove this result, but the proof is hidden  
 610 in the middle of a discussion culminating in the following  
 611 statement (simplified using our standing commutativity as-  
 612 sumption), to be compared with their earlier statement of  
 613 Theorem 5.1:

614 **Theorem 5.2.** *Any separated topological abelian group  $G$  is*  
 615 *isomorphic to a dense subgroup of a complete group  $\hat{G}$ . The*  
 616 *complete group  $\hat{G}$  (which is called the completion of  $G$ ) is*  
 617 *unique (up to isomorphism).*  
 618

619 In particular this theorem does not discuss at all whether  
 620 the hat notation denotes the same operation as in Theo-  
 621 rem 5.1. In this case it denotes the same *set*, and the am-  
 622 biguity about the uniform structure is not important since  
 623 they prove both constructions give the same structure. The  
 624 situation deteriorates once we notice the separation assump-  
 625 tion (which does not appear in Theorem 5.1). After proving  
 626 Theorem 5.2, Bourbaki writes<sup>5</sup>: Let  $G$  be a group which is not  
 627 necessarily separated, and let  $G' := G/\{0\}$  be the separated  
 628 group associated with  $G$ . The completion of  $G'$  is called the  
 629 separated completion of  $G$ , and denoted by  $\hat{G}$ . So we now  
 630 have a third meaning of the hat notation, still very closely  
 631 related, but definitely *not* the same set this time. Bourbaki  
 632 then states the universal property of  $\hat{G}$ , factoring morphisms  
 633 in the category of topological groups.

634 All this is pretty reasonable from a mathematical point  
 635 of view, because all three variants of the completion  $\hat{G}$  are  
 636 isomorphic. And we want the formalization to look just as  
 637 friendly as the real thing, so some thinking is needed. On top  
 638 of the fact that a formal definition needs to be one definition  
 639 rather than three definitions, there is a difficulty coming  
 640 from the type class mechanism. Whatever the definition  
 641 of completion  $G$ , Lean needs to find a uniform structure  
 642 instance, and a topological group instance, without enter-  
 643 ing an infinite loop, and without choosing an instance that  
 644 later lemmas won't automatically recognise as the right one.  
 645 Typically a lemma needing completeness of completion  $G$   
 646 could fail to derive this automatically if Lean infers a uniform  
 647 structure from the topological group structure, whether it  
 648 will definitely succeed if it gets it from the completion con-  
 649 struction (whose completeness is part of the conclusion of  
 650 Theorem 5.1).

651 We already explained part of the solution: using Theo-  
 652 rem 4.2 instead of Theorem 4.1, we can completely forget  
 653 about the separation assumption in Theorem 5.2. The only  
 654 trade-off is that we must give up the word subgroup since  
 655  $G \rightarrow \hat{G}$  will be injective only if  $G$  is separated. The other  
 656 part of the solution is to give up the idea that we have two  
 657 operations (building the uniform structure on a topological  
 658

659 <sup>5</sup>This is slightly edited for simplification purposes.

group and completing a uniform space) whose commutation should be proved and then silently used. Instead we introduce a predicate asserting compatibility of a uniform structure and a group structure. We then have a first lemma asserting that, from a topological group one can build a uniform structure whose underlying topology is definitionally equal to the original topology, and definitionally satisfies the compatibility predicate. And another lemma saying that if a group structure on  $G$  is compatible with a uniform structure  $\mathcal{U}$  then, on the completion of  $G$ , defined using only  $\mathcal{U}$ , the group structure obtained by extension is compatible with the completed uniform structure. Of course this last lemma has exactly the same non-trivial mathematical content as in the commutation assertion. But this repackaging avoids all the notation overloading and is formalization-friendly.

Once completions of abelian groups work nicely, the next task is to extend multiplication on the completion of a topological ring. As mentioned earlier, multiplication is not uniformly continuous, even in the elementary cases. So one must really check the limit assumption in Theorem 4.2, and the proof is pretty subtle. Bourbaki builds this extension in two steps because of the injectivity assumption hidden in 4.1. In particular the meaning of the hat notation is not quite the same as in their general discussion of completions. Our formalization does it in one step and proves, without silently changing the meaning of notations:

**Theorem 5.3.** *Let  $R$  be a topological ring, equipped with its natural uniform structure. There is a ring structure on the separated completion  $\hat{R}$  which is compatible with its topological structure, and such that  $\iota_R$  is a ring morphism. If  $f : R \rightarrow S$  is a continuous ring morphism between topological rings then  $\hat{f} : \hat{R} \rightarrow \hat{S}$  is also a ring morphism.*

The main point of this part of the story is that ordinary mathematics sets up topological ring completions by silently identifying different isomorphic mathematical objects, but being annoyed when trying to formalize led us to actually streamline the theory by proving Theorem 4.2 and introduce abstract completions.

Before realizing that Theorem 4.2 holds, we formalized Bourbaki's two step completion of topological rings, and encountered some formalization difficulty. Briefly, the general theory of uniform spaces give us a minimal equivalence relation with a separated quotient. For topological groups, one can prove  $x \sim y$  if and only if  $x - y$  is in the closure of the singleton zero. But this has mathematical content, and is certainly not definitional. So the topology library endows this quotient with a uniform structure, the algebraic theory endows it with a group structure, but these structures live on different quotient *types*. One can think of ways to solve that problem more elegantly than we did, but we didn't pursue our investigations in this direction since, as explained above, we found a way to completely bypass this issue.

## 5.4 Field completions

We also need topological fields, i.e. topological rings  $K$  where each non-zero has an inverse, and inversion is continuous from  $K^* := K \setminus \{0\}$  to itself. Complications arise because their completions, endowed with their ring structure built in the previous section, can have zero divisors. Bourbaki gives the following criterion, whose proof is basically left to the reader (the only hint being to use Theorem 4.1).

**Theorem 5.4** ([3, Proposition III.6.8.7]). *The completion of a separated topological field is a topological field if (and only if) the image under the mapping  $x \mapsto x^{-1}$  of every Cauchy filter which does not have a cluster point at zero, is a Cauchy filter.*

The separation assumption is not a restriction because a topological field is either separated or has the indiscrete topology. In the later case the separated completion is the zero ring, which is obviously not a field. In the formalization, the first question is how to handle the condition that inversion is continuous from  $K^*$  to itself. This definition implicitly equips  $K^*$  with the induced topology. Initially we used  $K^*$  as a subtype, mapping to  $K$  by inclusion and equipped with the induced topology, and induced uniform structure. This was manageable and we were able to prove the completion criterion, but every step was slightly painful, because inclusions are harder to ignore in formalized type theory.

On the algebraic side, it was decided long ago that manipulating inversion didn't deserve a subtype. Inversion is defined on  $K$ , and the definition of a field requires  $0^{-1} = 0$ . For the purpose of this discussion, let us denote by  $\text{inv}$  this weird function from  $K$  to itself. We can then require that, in a topological field,  $\text{inv}$  is continuous at every non-zero point, and this is equivalent to the previous definition. In the proof of Theorem 5.4, we apply the extension result, Theorem 4.2, to  $\iota_K \circ \text{inv}$ . Now the behavior of  $E_i(\iota_K \circ \text{inv})$  at zero is pretty mysterious, since it uses the axiom of choice to pick an element of  $\hat{K}$ . So we define  $\widehat{\text{inv}} : \hat{K} \rightarrow \hat{K}$  as mapping 0 to 0, and use  $E_i(\iota_K \circ \text{inv})$  elsewhere. Modulo this little trifle at 0, everything proceeds very smoothly. Because Theorem 4.2 is local, and  $K^*$  is a neighborhood of each of its elements (from the separation assumption), we don't need to care about zero when proving continuity of  $\widehat{\text{inv}}$  at non-zero points of  $\hat{K}$ . We defined a type class `completable_top_field` which records the assumptions of Theorem 5.4, in order for the type class system to automatically derive the field structures we will need. Of course it remains to provide the relevant instances of `completable_top_field`, but this is a story for the next section.

## 6 Valuations

### 6.1 Definition and target monoids

A totally ordered group (resp. totally ordered monoid) is a commutative group (resp. commutative monoid)  $\Gamma$  (whose composition law is written multiplicatively) together with

a total order such that the multiplication map  $\gamma' \mapsto \gamma\gamma'$  is order preserving for every  $\gamma$  in  $\Gamma$ . To every totally ordered group  $\Gamma$  one can associate a totally ordered monoid  $\Gamma_0 := \Gamma \cup \{0\}$  where the multiplication and order relation are extended by specifying that  $0\gamma = 0 \times 0 = 0$  and  $0 \leq \gamma$  for every  $\gamma$  in  $\Gamma$ .

A valuation on a ring  $R$  is a map  $v$  from  $R$  to  $\Gamma_0$  for some totally ordered group  $\Gamma$  such that:

- $v(0) = 0$  and  $v(1) = 1$
- $v(xy) = v(x)v(y)$  for all  $x$  and  $y$
- $v(x + y) \leq \max(v(x), v(y))$  for all  $x$  and  $y$ .

This terminology is a bit unfortunate, and not consistent with [4], but it is systematically used in the adic spaces literature, although a valuation in this sense is closer to a norm or a seminorm than a traditional valuation.

Notice how, once again, the above definition focuses on the construction of  $\Gamma_0$ , starting from an ordered group and adding an element, rather than the properties of the resulting object. The first consequence of this psychological trick is to help to hide the use of many formal properties of  $\Gamma_0$ . For instance we need at some point that, for every  $x, y$  and  $z$  in  $\Gamma_0$ ,  $xz < yz$  implies  $x < y$ . Such kind of results are not even stated in [4], and the systematic proof is extremely boring. There is a case split according to whether  $x, y$  and  $z$  belong to the original  $\Gamma$  or are 0, so there are eight possibilities, but only the case where all elements are non-zero is actually relevant, and needs a lemma about totally ordered groups. Of course our formalization cannot ignore these lemmas.

Initially we formalized this story using the type constructor `with_zero`  $\Gamma$  which is simply a wrapper for `option`  $\Gamma$  but has all the relevant type class instances (assuming  $\Gamma$  has its own relevant instances). We used Lean's coercion mechanism to automatically insert the "inclusion" map from  $\Gamma$  to  $\Gamma_0$  (in set theory, this is indeed a true inclusion, but there are no inclusions between types in type theory as a term can only have one type). The `norm_cast` tactic, by Paul-Nicolas Madeleine [18], greatly alleviates the pain of invoking lemmas about such coercions.

We then addressed the proof issue by taking advantage of how convenient it is to write specialized easy automation in Lean. Our tactic `with_zero_cases` made the case distinction and dispelled all irrelevant cases using Lean's `simplifier` and `norm_cast`, without any human assistance, leaving only the relevant case to prove by hand.

However that does not solve the following discrepancy. In an elementary context, valuations take value in  $\mathbb{R}_{\geq 0}$  or  $\mathbb{Q}_{\geq 0}$ . Set-theoretically, it is almost true that  $\mathbb{R}_{\geq 0} = (\mathbb{R}_{>0})_0$ . This tiny lie only requires that we forget that the 0 in  $(\mathbb{R}_{>0})_0$  is meant to be an extra element coming from nowhere in particular, hence has nothing to do with the neutral element of the additive group  $\mathbb{R}$ . It gets harder to ignore with type theoretic foundations. It means that the elementary theory with our abstract theory would require non-trivial glue. So we stepped back and gave up using `with_zero`. We defined

the concept of totally ordered commutative monoid with zero as a type equipped with a composition law, a total order, and special elements 0 and 1 with enough properties to guarantee that all instances are isomorphic (as ordered commutative monoids) to some  $\Gamma_0$ . This corresponds to a type class which admits both `with_zero`  $\Gamma$  and `{x :  $\mathbb{R}$  //  $0 \leq x$ }` as instances. Once again, focussing on properties instead of constructions gives us the needed extra flexibility at no cost.

## 6.2 Valuation topology and valuation extensions

In order to construct a valuation on stalks in Section 9, we will start with a field equipped with a valuation  $v$ , endow that field with a topology associated to  $v$ , complete, and extend  $v$  to the completion.

A valuation  $v$  on a ring  $R$  with values in  $\Gamma_0$  defines a topology characterized by the fact that, for every  $x$  in  $R$ , neighborhoods of  $x$  are subsets containing  $\{y \mid v(y - x) < \gamma\}$  for some  $\gamma \in \Gamma$  (existence of this topology needs to be proven of course). The first main result of this section is:

**Proposition 6.1.** *The topology coming from a valuation on a ring makes it a non-Archimedean topological ring. If this ring is (commutative and) a field, then it is completable, and the valuation extends to the completion, with the same target monoid.*

In this statement, non-Archimedean means having a fundamental system of neighborhood of zero consisting of additive subgroups. We now focus on the extension part of the proposition, for a field  $K$ . The elementary example is  $K = \mathbb{Q}$  equipped with some  $p$ -adic valuation, and the extension is the  $p$ -adic valuation on  $\mathbb{Q}_p$ .

The proof in [4, VI.5.3] is extremely short. Translated into our notation, it says: According to Proposition 1,  $v|_{K^*}$  is a continuous representation of  $K^*$  into  $\Gamma$  (equipped with the discrete topology), hence can be extended to a continuous representation  $\hat{v}$  of  $\hat{K}^*$  into  $\Gamma$ . The relation  $\hat{v}(x + y) \leq \max(\hat{v}(x), \hat{v}(y))$  holds in  $K^*$ , hence stays true in  $\hat{K}^*$  by continuity. Hence  $\hat{v}$ , extended by  $\hat{v}(0) = 0$ , is a valuation.

At first reading, we thought the first "hence" was implicitly referring to the extension promised by Theorem 5.1. But this would require identifying  $\hat{K}^*$  with the uniform completion of the multiplicative group  $K^*$ . However the uniform structure induced by  $K$  on  $K^*$  is not the uniform structure associated to the topological group  $(K^*, \times)$ , otherwise multiplication would be uniformly continuous. After exchanging a couple of emails with current and former collaborators of Nicolas Bourbaki, it looks like Bourbaki meant to apply [3, III.3.3.5]. In the meantime, we formalized another proof.

Before sketching this alternative proof, let us consider the non-Archimedean triangle inequality argument in the above proof. It looks rather uncontroversial until one notices that  $x + y$  can land outside  $K^*$  (or  $\hat{K}^*$ ) even if  $x$  and  $y$  are both in  $K^*$  (or  $\hat{K}^*$ ). This is a bit embarrassing, but not a serious difficulty: one can separately argue that if  $x + y = 0$  in  $\hat{K}$



then the inequality holds for free. But this will complicate the formal proof, which also requires spelling out other details.

Fortunately, all this is avoided by our alternative proof. We endow  $\Gamma_0$  with a topology where non-zero points are open but neighborhoods of zero are subsets containing some open interval  $\{\gamma \mid \gamma < \gamma_0\}$  (this is not an exotic topology to put on  $\Gamma_0$ , but it seems Bourbaki tried to avoid introducing it). We then apply Theorem 4.2 to the full map  $v : K \rightarrow \Gamma_0$ . Proving the assumptions of the theorem involves, as in the elementary proofs, arguing that the valuation is locally constant on  $K$ . Of course we also need to prove that  $\Gamma_0$  is a regular topological space and its order relation plays nicely with the topology.

The conclusion is always the same: in order to avoid formalization pain when filling in details or fixing little mistakes, we used a slightly different point of view which turns out to be more natural. Indeed one cannot extend a valuation by seeing it purely as a multiplicative group morphism and hope that it will play nicely with the triangle inequality, which is about addition.

## 7 Presheaves and sheaves

A presheaf  $\mathcal{F}$  on a topological space  $X$  with values in a category  $\mathcal{C}$  associates an object  $\mathcal{F}(U)$  of  $\mathcal{C}$  to each open subset  $U$  of  $X$ , and a so-called restriction morphism  $\text{res}_{U,V} : \mathcal{F}(U) \rightarrow \mathcal{F}(V)$  whenever  $V \subseteq U$ , with  $\text{res}_{U,W} = \text{res}_{U,V} \circ \text{res}_{V,W}$  when  $W \subseteq V \subseteq U$ . When  $\mathcal{C}$  is a concrete category, as in all our examples, elements of  $\mathcal{F}(U)$  are called sections of  $\mathcal{F}$  on  $U$ . The most elementary examples are sheaves of functions, for instance  $\mathcal{F}(U)$  could be the ring of real-valued continuous functions on  $U$ , with the usual restriction operation. In our context, the structure sheaves of adic spaces will be thought as sheaves of analytic functions, but their definition will involve nothing like functions (see Theorem 8.1 below). In particular building restriction morphisms, and checking their compatibility, will involve abstract work.

In order to reasonably think of sections of a presheaf as functions, they must satisfy some locality and gluing conditions. One then says that  $\mathcal{F}$  is a sheaf. Roughly, for each open covering  $(U_i)$  of an open set  $U$ , and each collection of sections  $s_i \in \mathcal{F}(U_i)$  that are compatible, i.e.  $\text{res}_{U_i, U_i \cap U_j} s_i = \text{res}_{U_j, U_i \cap U_j} s_j$  for all  $i$  and  $j$ , there is a unique  $s \in \mathcal{F}(U)$  which restricts to  $s_i$  on each  $U_i$ . The precise definition more carefully uses the full structure of  $\mathcal{C}$ . In our case,  $\mathcal{C}$  is the category of complete separated topological rings, and the bijection between collections of compatible sections  $(s_i)$  and  $s$  must be a homeomorphism.

As is often the case, we will construct sheaves by first defining them on elements of a basis  $\mathcal{B}$  of open subsets, and then we extend by setting  $\mathcal{F}(U) = \varprojlim_V \mathcal{F}(V)$  the projective limit being over elements of  $\mathcal{B}$  containing  $U$ . In practice the partially ordered set over which we took the projective limit was slightly finer than the obvious one, however one can

prove that the limit we took is naturally isomorphic to the “correct” limit. Because projective limits are only defined up to isomorphism, this is not a problem.

We will also need local information encoded in stalks of  $\mathcal{F}$  at a point  $x$ ,  $\mathcal{F}_x = \varinjlim_{U \ni x} \mathcal{F}(U)$ , where the inductive limit is over open subsets containing  $x$ . When the sheaf is constructed from a basis, one can prove it is sufficient to consider basic subsets containing  $x$ . In our situation, we will first forget about the topologies on  $\mathcal{F}(U)$  and consider stalks of sheaves of rings.

Sheaves of rings were introduced in Lean during the formalization of schemes [5]. We reused that formalization, extending it to sheaves of *topological* rings. This is an ad hoc formalization which does not rely on mathlib’s category theory library. We found this to be easier for our purposes, but it is not sustainable since turning to sheaves with values in other categories would need a lot of code duplication.

## 8 Adic spectra

Everything in Sections 4 to 6 was already present in [3, 4] going back to the 40’s and 60’s respectively. Sheaves of topological rings are also from the same era. We now jump to the 90s with the construction of  $\text{Spa}(A, A^+)$ , the adic spectrum of a Huber pair.

### 8.1 Valuation spectra and continuous valuations

The first step towards constructing  $\text{Spa}$  is to consider the valuation spectrum  $\text{Spv}(R)$  of a ring  $R$ . The informal definition starts by saying that two valuations  $v : R \rightarrow \Gamma_0$  and  $w : R \rightarrow \Delta_0$  are *equivalent* if there exists an isomorphism  $\varphi$  between the monoids  $v(R)$  and  $w(R)$  such that  $w = \varphi \circ v$ . Then  $\text{Spv}(R)$  is defined as the “set” of “equivalence classes” of valuations on  $R$ . But there is no set of valuations, because the target monoids can be arbitrarily large. In type theory, this translates into universe issues. Fortunately there are well-known reformulations that avoid these issues. In particular, two valuations are equivalent if and only if they pull back the order relation on their target monoids to the same preorder on  $R$ . Of course preorders on  $R$  form a set, and one can define  $\text{Spv}(R)$  as the subset of those coming from a valuation. Since we still want to pretend  $\text{Spv}$  is defined as a quotient, we then prove a series of lemmas which provides the usual quotient interface.

The next construction starts with a topological ring  $R$ . Here there is a notion of a continuous valuation  $v : R \rightarrow \Gamma_0$ , which is more subtle than asking for continuity when the target is equipped with the topology discussed in the previous section. It goes through replacing the valuation by some canonical representative of its “equivalence class” before checking continuity (one for which  $\Gamma$  is as small as possible). This enables us to define the subtype  $\text{Cont}(R) \subseteq \text{Spv}(R)$  of “continuous valuations” on  $R$ , where the quotation marks highlight both the fact that equivalence classes of

valuations are now called valuations, and that they are not even equivalence classes.

## 8.2 $\text{Spa}(A, A^+)$ and its structure presheaf

In discrete commutative algebra a large role is played by Noetherian rings, because of their finiteness properties. In the theory of adic spaces, there are no Noetherian assumptions. However, the rings that show up are all endowed with a topology, and one can recover finiteness constraints by asking for a subtle interplay between the topological and algebraic structures. *Huber rings* form such a class of topological rings.

A topological ring  $A$  is Huber if there exists some open subring  $A_0$  whose topology is  $I$ -adic for some finitely generated ideal  $I$ . A Huber pair  $(A, A^+)$  is a Huber ring equipped with an open subring which is integrally closed and power bounded (the definition of those conditions will not be relevant below). As a sanity check for our formalization, we proved that  $(\mathbb{Q}_p, \mathbb{Z}_p)$  is a Huber pair. Doing that already reveals that all those definitions involving open subrings are stated too rigidly for type theory. Indeed `mathlib` defines  $\mathbb{Z}_p$  as a subtype, in order to equip it with all its extra structures through the type class mechanism. So we replaced open subrings by open embeddings of rings.

As a set,  $\text{Spa}(A, A^+) = \{v \in \text{Cont}(A) \mid \forall x \in A^+, v(x) \leq 1\}$ . We will now set  $X = \text{Spa}(A, A^+)$  for some fixed Huber pair. In order to put a topology on  $X$ , to each  $s$  in  $A$  and each finite subset  $T \subseteq A$  such that  $T \cdot A$  is open, one associates the so-called rational open subset  $R(T/s) := \{v \in X \mid v(s) \neq 0 \wedge \forall t \in T, v(t) \leq v(s)\}$ . Here  $T \cdot A$  is the additive subgroup generated by products of elements of  $T$  and general elements of  $A$ . We also associate to each valuation  $v$  on  $A$  its support  $\text{supp}(v) = \{x \in A \mid v(x) = 0\}$ . This is a prime ideal, hence  $A/\text{supp}(v)$  has a field of fractions  $K_v$ . The valuation  $v$  extends to  $K_v$  where it defines a topological field structure as in Section 6.2.

**Theorem 8.1.** *Let  $(A, A^+)$  be a Huber pair, and  $X = \text{Spa}(A, A^+)$ .*

1. *The rational open subsets  $R(T/s)$  form a topology basis on  $X$ .*
2. *For each  $(T, s)$  as above, there is a topology on the localization  $A[s^{-1}]$  turning it into a non-Archimedean topological ring, denoted by  $A(T/s)$ , which is universal for the following properties: the localization map  $\varphi : A \rightarrow A(T/s)$  is continuous,  $\varphi(s)$  is invertible, and the image  $(t \mapsto \varphi(t)\varphi(s)^{-1})(T)$  is power-bounded.*
3. *There is a presheaf  $\mathcal{O}_X$  of complete separated topological rings on  $X$  whose value on each  $R(T/s)$  is naturally isomorphic to the separated completion  $\widehat{A\langle T/S \rangle}$  of the above ring.*
4. *For every  $x$  in  $X$ , there is an equivalence class  $v_x$  of valuations on the stalk  $\mathcal{O}_{X,x}$  such that, for every  $v$  representing  $x$ , the representatives of  $v_x$  factor through  $\widehat{K}_v$ .*

Explaining the proof of this theorem is far beyond the scope of this paper, but we want to focus on some issues

that arose during its formalization, and also indicate how the previous sections come together in the proof. This will be done in Sections 8.3 to 8.5 below.

Before doing that, we need a disclaimer. An important theorem in the theory, necessary for any significant further work, states that two natural topologies on  $X$  coincide; the topology on  $X$  is thus defined by mathematicians to be “both of them”. The proof that the topologies coincide is Theorem 3.5(ii) of [14] or Theorem 7.35(2) of [25]. We did *not* formalise the proof of this theorem. The formal proof would probably involve months of work, developing a good interface for primary and secondary specialisations of valuations on rings, as is done in Section 2 of [14] and its references, or sections 4 and 7 of [25]. The topology described above is the best suited to our needs, although this shortcut creates some additional technical issues. In short, our proof that our definition of an adic space agrees with “the definition in the literature” invokes a theorem in the literature which we have not yet formalised.

## 8.3 Localization, elaboration, and the community

The first issue we want to discuss is that of balancing readable statements against the demands of Lean’s elaborator. A typical lemma that shows up in the proof of the second item of Theorem 8.1 is the following .

**Lemma 8.2.** *Let  $A$  be a Huber ring and let  $T$  be a subset of  $A$  such that ideal generated by  $T$  is open. For every open subgroup  $U$  of  $A$ ,  $T \cdot U$  is open.*

Our formalisation of this statement reads as follows:

```
lemma mul_T_open {A : Type*} [Huber_ring A]
  {T : set A} (hT : is_open ((ideal.span T) : set A))
  (U : open_add_subgroup A) :
  is_open (↑(T · span ℤ (U : set A)) : set A) :=
```

This statement features several type ascriptions, an explicit coercion operator  $\uparrow$ , and an ad-hoc coercion trick. The coercion mechanism in Lean 3 works great in simple situations, but not always when several instances must be resolved in the correct order. In `mathlib`, the type `ideal A` bundles a subset of  $A$  with the expected properties. It has a coercion to `set A`. But in assumption `hT`, Lean also needs to figure out the topological space that `is_open` refers to. The type ascription helps it doing those things in the correct order. Things pile up even more in the conclusion of the lemma, where the (bundled) open subgroup  $U$  is seen as a  $\mathbb{Z}$ -submodule by first forgetting everything but the underlying subset and then taking the  $\mathbb{Z}$ -span to build a (bundled) submodule whose underlying set is exactly the same. Lean then needs to figure out the meaning of  $\cdot$  which is the overloaded notation for scalar multiplication (more on that instance below), and then again find the relevant topological space. Coercions to subsets are especially tricky because `set X` is defined as the function type  $X \rightarrow \text{Prop}$ , and coercions to functions receive

a special treatment, which can be misleading in the set case. Using the explicit coercion operator helps because coercions to functions have a different dedicated operator.

In conclusion, we managed to write a somewhat recognizable statement, but the situation is not perfect. It may be improvable by more expert Lean users, but we still claim this is not as easy as we'd like it to be.

The second issue is related to the first, and has to do with the notation  $T \cdot U$ . One of the benefits of introducing this notation (and the corresponding instances of algebraic type-classes) is that it allows to use all sorts of results from the algebraic library: submodules of an algebra form a semiring, that is a semialgebra of the semiring of subsets (under elementwise multiplication, followed by the linear span).

However, not all of these facts were available in `mathlib` at the time of our development. The following instances were available (here  $A$  is an  $R$ -algebra): `semigroup (submodule R A)`, `comm_semiring (ideal R)`, and `algebra R R`.

In our project, we needed `semiring (submodule R A)`. However, when we added this instance to the system, type class search would be able to find to different instances of `semiring (ideal R)`, and they were not definitionally equal. One came from the instance of `comm_semiring (ideal R)` in `mathlib`. The other came from combining the two instances `semiring (submodule R R)` and `algebra R R`, since `ideal R` is defined to be `submodule R R`.

The solution, of course, was to change `mathlib`, generalizing the `semigroup (submodule R A)` to a `semiring` and using this new instance to redefine the existing instance of `comm_semiring (ideal R)`. This change was proposed in pull request #856 to `mathlib`. As long as this pull request was not merged upstream, making progress on localisations of Huber rings was not really possible. This very unfamiliar experience for mathematicians was frustrating, although it lasted only one week.

At the time the maintainers of `mathlib` had very little time for review and merging of pull requests. Altogether this episode fostered reflection about the processes involved in maintaining `mathlib`. Our overall experience is that maintenance is now smoother than before. Nevertheless, this experience raises questions for the future. Are lock-ups like the one explained above expected to occur more often when libraries and formalisation efforts scale up?

## 8.4 The structure presheaf

We now comment on the third item of Theorem 8.1. This was more problematic than it should have been, because of our incomplete interface to the topological space structure on  $X$ . Recall that if  $s \in A$  and  $T \subseteq A$  is a finite subset such that  $T \cdot A$  is open, then there is a rational open subset  $R(T/s)$  of  $X$  defined by some inequalities. We would like the ring of functions on  $R(T/s)$  to be the completion  $A\langle T/s \rangle$  of  $A(T/s)$ . Because the rational open subsets form a basis for the topology (by Theorem 8.1(1)), a function on a general open set  $U$

should be determined by its restrictions to the rational open subsets contained in  $U$ . So we would like to define  $\mathcal{O}_X(U)$ , for  $U$  a general open subset of  $X$ , to be the subset of the product of the  $A\langle T/s \rangle$  such that  $R(T/s) \subseteq U$ , consisting of functions which “agree on overlaps”. However it is rather a deep result about continuous valuations which shows that if  $R(T_1/s_1) \subseteq R(T_2/s_2)$  is an inclusion of rational open subsets, then there is a natural ring homomorphism  $A\langle T_2/s_2 \rangle \rightarrow A\langle T_1/s_1 \rangle$ , which is an isomorphism if the inclusion is the identity, and we have not yet formalised this result. We avoided this issue by proving that  $R(T_1/s_1) \cap R(T_2/s_2) = R(T_1T_2/s_1s_2)$  and defining ring homomorphisms  $A\langle T_1/s_1 \rangle \rightarrow A\langle T_1T_2/s_1s_2 \rangle$  and  $A\langle T_2/s_2 \rangle \rightarrow A\langle T_1T_2/s_1s_2 \rangle$ , and defining  $\mathcal{O}_X(U)$  to be the elements  $(f_i)$  of the product of the  $A\langle T_i/s_i \rangle$  for which  $R(T_i/s_i) \subseteq U$ , such that the restrictions of  $f_i$  and  $f_j$  agreed in  $A\langle T_iT_j \rangle/s_1s_2$ . This definition is mathematically equivalent to the usual one, although the obstruction to formalising the proof of this in Lean is that we need primary and secondary specialisations of valuations, which we do not yet have. However it is not too difficult to show in Lean that our definition satisfies the axioms of a presheaf, which is what we need.

## 8.5 Stalk valuations

Having defined the presheaf  $\mathcal{O}_X$  on  $X$ , the last piece of structure we need is a valuation on each stalk. For a point  $x$  of  $X$ , let  $v$  be a valuation in its equivalence class. The stalk of  $\mathcal{O}_X$  at  $x$  is the colimit of the rings  $\mathcal{O}_X(U)$  as  $U$  runs through the open subsets of  $X$  containing  $x$ . Because the rational open subsets  $R(T/s)$  form a basis for the topology, this colimit is isomorphic to the colimit of the rings  $A\langle T/s \rangle$  where  $R(T/s)$  runs through the rational open subsets of  $X$  containing  $x$ , so our task is to put a valuation on this colimit which naturally extends  $v$ . The universal property of the colimit tells us that we can define a ring homomorphism from the colimit by giving a collection of ring homomorphism from each  $A\langle T/s \rangle$  which are compatible with our restriction maps. The valuation  $v$  extends to a valuation  $\hat{v}$  on  $K_v = \text{Frac}(A/\text{supp}(v))$  and, by Section 6.2, to a valuation on its completion. Thus it suffices to give ring homomorphisms  $A\langle T/s \rangle \rightarrow \widehat{K}_v$  for each pair  $(T, s)$  such that  $x \in R(T/s)$ , which are compatible with all restriction maps. The assumption  $x \in R(T/s)$  implies that  $s \notin \text{supp}(v)$ . Hence the map  $A \rightarrow K_v$  extends to a ring homomorphism  $A[s^{-1}] \rightarrow K_v$ . The fact that  $x \in R(T/s)$  also implies that this homomorphism is continuous on  $A(T/s)$  which is  $A[s^{-1}]$  endowed with the topology coming from  $T$ , and we deduce the existence of a ring homomorphism  $A\langle T/s \rangle \rightarrow \widehat{K}_v$  using the completion functor. It is formal to check that these ring homomorphisms are compatible with our restriction maps and hence they give rise to a ring homomorphism from the stalk of  $\mathcal{O}_X$  at  $x$  to  $\widehat{K}_v$ ; composing with the valuation  $\hat{v}$  on  $\widehat{K}_v$  gives us the valuation we require.

In conclusion, to a Huber pair  $(A, A^+)$  we are able to associate its adic spectrum  $X = \text{Spa}(A, A^+)$ , together with a presheaf of topological rings, and a valuation on each stalk of the presheaf.

## 9 Adic and perfectoid spaces

An adic space is a topological space equipped with a sheaf of complete topological rings with valuations on stalks which is locally isomorphic to the adic spectrum of a Huber pair; see below for a formal definition. These isomorphisms are taking place in an auxiliary category which only plays a role in the definition of an adic space; this category does not seem to be taken too seriously by mathematicians and it does not even seem to have an established name. References [14] and [25] call it  $\mathcal{V}$ , and we call it CLVRS (for “complete locally valued ringed space”). Its objects consist of the following data: a topological space  $X$ , a sheaf of complete topological rings  $\mathcal{O}_X$  on  $X$ , and for each  $x \in X$  an equivalence class of valuations  $v_x$  on the stalk of  $\mathcal{O}_X$  at  $x$ , whose support is the unique maximal ideal of  $\mathcal{O}_{X,x}$ . It is clear, although rarely explicitly mentioned, that an open subspace of a topological space equipped with such structure, naturally inherits such a structure. We adopt the usual mathematical tradition of conflating the object  $(X, \mathcal{O}_X, (v_x)_{x \in X})$  of  $\mathcal{V}$  with its underlying topological space  $X$ . A morphism from the object  $(X, \mathcal{O}_X, (v_x)_{x \in X})$  to  $(Y, \mathcal{O}_Y, (v_y)_{y \in Y})$  consists of a continuous map  $f : X \rightarrow Y$ , and a morphism of sheaves of topological rings  $\mathcal{O}_Y \rightarrow f_*\mathcal{O}_X$  such that  $v_x$  pulls back to  $v_{f(x)}$  along the induced morphism  $\mathcal{O}_{Y,f(x)} \rightarrow \mathcal{O}_{X,x}$  of stalks. This gives us a notion of isomorphism in this category.

In [25] we see a larger category  $\mathcal{V}^{\text{pre}}$  where  $\mathcal{O}_X$  is only assumed to be a presheaf rather than a sheaf. We used an even more general category `PreValuedRingedSpace`, and our a definition of an adic space is an object of CLVRS which is locally isomorphic (in `PreValuedRingedSpace`) to the continuous spectrum of a Huber pair:

```
def is_adic_space (X : CLVRS) :=
  ∀ x, ∃ (U : opens X) (A : Huber_pair), x ∈ U ∧ (Spa A ≅ U)
```

Since CLVRS is a full subcategory of `PreValuedRingedSpace`, this is equivalent to the usual definition. So the last main piece of work is to set up those two categories. Instantiating  $\text{Spa}(A, A^+)$  as an object of `PreValuedRingedSpace` is mostly restating things we did before.

Once the definition of an adic space is formalised, it is not hard to formalise the definition of a perfectoid space; it simply modifies the previous definition by asking that each Huber pair is perfectoid. Being perfectoid is just a relatively straightforward predicate on a Huber pair, demanding essentially that there is a prime number  $p$  and an open subring  $A_0$  of  $A$  such that  $x \mapsto x^p$  is a surjection from  $A_0/p$  to itself, together with the “deep ramification” condition mentioned in the introduction. This condition forces every element of  $A_0$  to have a  $p$ th root modulo  $p$ ; this algebraic predicate implies

that many functions on the space have  $p$ th roots which are also functions on the space, which has non-trivial geometric implications. Scholze’s insight that this is an important condition cannot be explained here, although its importance in modern mathematics cannot be understated.

However, we have almost nothing beyond the definition. We have one example – the empty perfectoid space. Even this example involved some work, boiling down to the statement that the unique map from an infinite product of trivial topological rings to the trivial topological ring was an isomorphism; this is what the sheaf axiom boils down to in this case. We are close to giving examples where the underlying space has one element, but formalising non-trivial examples will be harder.

## 10 Future work and concluding thoughts

We have developed a robust theory of completion of topological groups and rings in the Lean theorem prover. We have proved universal properties which characterise our constructions. We have tested the usability of our theory by developing a theory of continuous valuations and adic spectra on top of it (and enough sheaf theory). We developed the theory of adic spectra far enough to be able to formalise the definitions of adic and perfectoid spaces. Very natural future work would be to develop the theory of primary and secondary specialisations of valuations, which would allow to start developing properties of adic and perfectoid spaces.

We feel that, since a proof assistant can handle the definition of a perfectoid space, then they should be able to handle a general modern mathematical definition. Such modern definitions are uncommon in the formalisation world, perhaps because so few mathematicians know how to use formalisation software. This is a sad state of affairs which we hope will change. Hopefully our contribution will help a bit clearing the misconception that proof assistants are only usable by people with a computer science degree, and only for proving things about combinatorics [10] or finite group theory [11]. We already received numerous invitations to talk about this project in mathematics departments. The Notices of the AMS, and their British and French analogues all asked us to write surveys about our experience as mathematicians using a proof assistant. So we can claim that at least some mathematicians noticed our work. On the other side of the bridge, we hope our efforts will help encouraging the development of proof assistants to make them easier to use by mathematicians.

## References

- [1] Anonymous. 2019. The Lean mathematical library. <https://leanprover-community.github.io/papers/mathlib-paper.pdf>
- [2] Sylvie Boldo, Catherine Lelay, and Guillaume Melquiond. 2015. Coquelicot: A User-Friendly Library of Real Analysis for Coq. *Mathematics in Computer Science* 9, 1 (01 Mar 2015), 41–62. <https://doi.org/10.1007/s11786-014-0181-1>

- 1321 [3] Nicolas Bourbaki. 1966. *Elements of mathematics. General topology.*  
1322 *Part 1.* Hermann, Paris; Addison-Wesley Publishing Co., Reading,  
1323 Mass.-London-Don Mills, Ont. vii+437 pages.
- 1324 [4] Nicolas Bourbaki. 1998. *Commutative algebra. Chapters 1–7.* Springer-  
1325 Verlag, Berlin. xxiv+625 pages. Translated from the French, Reprint  
1326 of the 1989 English translation.
- 1327 [5] Kevin Buzzard, Ramon Fernández, Chris Hugues, and Lau Kenny. 2019.  
1328 Schemes in Lean. <https://github.com/ramonfmir/lean-scheme>
- 1329 [6] Thierry Coquand and Christine Paulin. 1988. Inductively defined types.  
1330 In *COLOG-88, International Conference on Computer Logic, Tallinn, USSR, December 1988, Proceedings.* 50–66. [https://doi.org/10.1007/3-540-52335-9\\_47](https://doi.org/10.1007/3-540-52335-9_47)
- 1331 [7] Sander R. Dahmen, Johannes Hölzl, and Robert Y. Lewis. 2019. For-  
1332 malizing the Solution to the Cap Set Problem. *CoRR* abs/1907.01449  
1333 (2019). arXiv:1907.01449 <http://arxiv.org/abs/1907.01449>
- 1334 [8] Leonardo Mendonça de Moura, Soonho Kong, Jeremy Avigad, Floris  
1335 van Doorn, and Jakob von Raumer. 2015. The Lean Theorem Prover  
1336 (System Description). In *Automated Deduction - CADE-25 - 25th International Conference on Automated Deduction, Berlin, Germany, August 1-7, 2015, Proceedings.* 378–388. [https://doi.org/10.1007/978-3-319-21401-6\\_26](https://doi.org/10.1007/978-3-319-21401-6_26)
- 1337 [9] François Garillot, Georges Gonthier, Assia Mahboubi, and Laurence  
1338 Rideau. 2009. Packaging Mathematical Structures. In *Theorem Proving in Higher Order Logics, 22nd International Conference, TPHOLs 2009, Munich, Germany, August 17-20, 2009. Proceedings.* 327–342. [https://doi.org/10.1007/978-3-642-03359-9\\_23](https://doi.org/10.1007/978-3-642-03359-9_23)
- 1339 [10] Georges Gonthier. 2007. The Four Colour Theorem: Engineering of a  
1340 Formal Proof. In *Computer Mathematics, 8th Asian Symposium, ASCM 2007, Singapore, December 15-17, 2007. Revised and Invited Papers.* 333. [https://doi.org/10.1007/978-3-540-87827-8\\_28](https://doi.org/10.1007/978-3-540-87827-8_28)
- 1341 [11] Georges Gonthier, Andrea Asperti, Jeremy Avigad, Yves Bertot, Cyril  
1342 Cohen, François Garillot, Stéphane Le Roux, Assia Mahboubi, Rus-  
1343 sell O’Connor, Sidi Ould Biha, Ioana Pasca, Laurence Rideau, Alexey  
1344 Solovyev, Enrico Tassi, and Laurent Théry. 2013. A Machine-Checked  
1345 Proof of the Odd Order Theorem. In *Interactive Theorem Proving - 4th International Conference, ITP 2013, Rennes, France, July 22-26, 2013. Proceedings.* 163–179. [https://doi.org/10.1007/978-3-642-39634-2\\_14](https://doi.org/10.1007/978-3-642-39634-2_14)
- 1346 [12] Sébastien Gouëzel and Vladimir Shchur. 2019. A corrected quantitative  
1347 version of the Morse lemma. *Journal of Functional Analysis* 277, 4  
1348 (2019), 1258 – 1268. <https://doi.org/10.1016/j.jfa.2019.02.021>
- 1349 [13] Johannes Hölzl, Fabian Immler, and Brian Huffman. 2013. Type Classes  
1350 and Filters for Mathematical Analysis in Isabelle/HOL. In *Interactive Theorem Proving - 4th International Conference, ITP 2013, Rennes, France, July 22-26, 2013. Proceedings.* 279–294. [https://doi.org/10.1007/978-3-642-39634-2\\_21](https://doi.org/10.1007/978-3-642-39634-2_21)
- 1351 [14] Roland Huber. 1993. Continuous valuations. *Math. Z.* 212, 3 (1993),  
1352 455–477. <https://doi.org/10.1007/BF02571668>
- 1353 [15] Roland Huber. 1996. *Étale cohomology of rigid analytic varieties and adic spaces.* Friedr. Vieweg & Sohn, Braunschweig. x+450 pages. <https://doi.org/10.1007/978-3-663-09991-8>
- 1354 [16] Ioan James. 1999. *Topologies and uniformities.* Springer-Verlag  
1355 London, Ltd., London. xvi+230 pages. <https://doi.org/10.1007/978-1-4471-3994-2> Revised version of it Topological and uniform  
1356 spaces [Springer, New York, 1987; MR0884154 (89b:54001)].
- 1357 [17] Robert Y. Lewis. 2019. A formal proof of Hensel’s lemma over the  
1358  $p$ -adic integers. In *Proceedings of the 8th ACM SIGPLAN International Conference on Certified Programs and Proofs, CPP 2019, Cascais, Portugal, January 14-15, 2019.* 15–26. <https://doi.org/10.1145/3293880.3294089>
- 1359 [18] Paul-Nicolas Madeleine. 2019. Arithmetic and Casting in  
1360 Lean. [https://lean-forward.github.io/internships/arithmetic\\_and\\_](https://lean-forward.github.io/internships/arithmetic_and_casting_in_lean.pdf)  
1361 [casting\\_in\\_lean.pdf](https://lean-forward.github.io/internships/arithmetic_and_casting_in_lean.pdf) Internship report.
- 1362 [19] Assia Mahboubi and Enrico Tassi. 2017. Mathematical Components.  
1363
- 1364 [20] Álvaro Pelayo, Vladimir Voevodsky, and Michael A. Warren. 2015. A  
1365 univalent formalization of the  $p$ -adic numbers. *Mathematical Structures*  
1366 *in Computer Science* 25, 5 (2015), 1147–1171. <https://doi.org/10.1017/S0960129514000541>
- 1367 [21] Michael Rapoport. 2019. The work of Peter Scholze. (2019).  
1368 arXiv:1900.07222 <http://arxiv.org/abs/1909.07222>
- 1369 [22] Damien Rouhling. 2019. Outils pour la formalisation en analyse clas-  
1370 sique. [http://www-sop.inria.fr/members/Damien.Rouhling/data/phd/](http://www-sop.inria.fr/members/Damien.Rouhling/data/phd/thesis.pdf)  
1371 [thesis.pdf](http://www-sop.inria.fr/members/Damien.Rouhling/data/phd/thesis.pdf) PhD thesis draft.
- 1372 [23] Peter Scholze. 2012. Perfectoid spaces. *Publ. Math. Inst. Hautes Études*  
1373 *Sci.* 116 (2012), 245–313. <https://doi.org/10.1007/s10240-012-0042-x>
- 1374 [24] Peter Scholze. 2017. Étale cohomology of diamonds.  
1375 arXiv:arXiv:1709.07343
- 1376 [25] Torsten Wedhorn. 2012. Adic Spaces. arXiv:1910.05934
- 1377 [26] Torsten Wedhorn. 2019. On the work of Peter Scholze.  
1378 arXiv:1909.11510